

# Space of chord-arc curves and BMO/VMO Teichmüller space

KATSUHIKO MATSUZAKI and HUAYING WEI

**Abstract.** This paper focuses on the structure of the subspace  $T_c$  of the BMO Teichmüller space  $T_b$  corresponding to chord-arc curves, which contains the VMO Teichmüller space  $T_v$ . We prove that  $T_c$  is not a subgroup with respect to the group structure of  $T_b$ , but it is preserved under the inverse operation and the left and the right translations by any element of  $T_v$ . Moreover, we show that  $T_b$  has a fiber structure induced by  $T_v$ , and the complex structure of  $T_b$  can be projected down to the quotient space  $T_v \backslash T_b$ . Then, we see that  $T_c$  consists of fibers of this projection, and its quotient space also has the induced complex structure.

## Jännekaarikäyrien avaruus ja Teichmüllerin BMO/VMO-avaruus

**Tiivistelmä.** Tämä työ keskittyy Teichmüllerin BMO-avaruuden  $T_b$  jännekaarikäyriin liittyvän aliavaruuden  $T_c$  rakenteeseen. Tämä  $T_c$  sisältää Teichmüllerin VMO-avaruuden  $T_v$ . Osoitamme, että  $T_c$  ei ole aliryhmä avaruuden  $T_b$  ryhmärakenteen suhteen, mutta  $T_c$  on kuitenkin suljettu sekä käänteisoperaation että aliavaruuden  $T_v$  suuntaisten vasemman- ja oikeanpuoleisten siirtojen suhteen. Lisäksi osoitamme, että  $T_v$  antaa avaruudelle  $T_b$  säierakenteen, ja että avaruuden  $T_b$  kompleksinen rakenne voidaan projisoida tekijäavaruuteen  $T_v \backslash T_b$ . Sen jälkeen näemme, että  $T_c$  koostuu tämän projektion säikeistä, ja että myös sen tekijäavaruudella on johdettu kompleksinen rakenne.

## 1. Introduction

Chord-arc curves have been studied with great interest in the fields of harmonic analysis and complex analysis. This concept is related to BMO and VMO functions. Astala and Zinsmeister [3] brought chord-arc curves and BMO functions into the theory of Teichmüller spaces and introduced the BMO Teichmüller space  $T_b$  as an analogous space with the universal Teichmüller space  $T$ . A brief summary concerning the background on this classical Teichmüller space of quasicircles and quasiconformal maps is given in Section 2. In this paper, we consider the structure of the space of chord-arc curves. We realize this space  $T_c$  in the BMO Teichmüller space  $T_b$  so that it contains the VMO Teichmüller space  $T_v$ . The precise definitions of these spaces are given in Section 3. The main theme of this paper is the investigation of the fiber structure of  $T_c$  with respect to the projection of  $T_b$  onto  $T_v$ .

In the universal Teichmüller space  $T$ , the little subspace  $T_0$  is defined by a certain vanishing property of its elements, and the quotient  $T_0 \backslash T$  was considered by Gardiner and Sullivan [13]. By taking the projection of the Bers embedding providing

---

<https://doi.org/10.54330/afm.122367>

2020 Mathematics Subject Classification: Primary 30C62, 30F60, 30H35; Secondary 28A75, 42A45, 58D05.

Key words: Chord-arc curve, BMO Teichmüller space, VMO Teichmüller space, strongly quasisymmetric, strongly symmetric, Carleson measure, quotient Bers embedding, asymptotic Teichmüller space.

Research supported by Japan Society for the Promotion of Science (KAKENHI 18H01125 and 21F20027) and by the National Natural Science Foundation of China (Grant No. 12271218).

© 2023 The Finnish Mathematical Society

a complex Banach manifold structure for  $T$  and its closed submanifold structure for  $T_0$ , they showed that the asymptotic Teichmüller space  $AT = T_0 \backslash T$  possesses the quotient Banach manifold structure. A formulation of this result in terms of the foliation of  $T$  by  $T_0$  is given in Theorem A and Corollary B. Our first result stated in Theorem 1 is an analogue of this formulation in the relation between  $T_b$  and  $T_v$ , which is based on Theorem C about the Bers embedding and the complex Banach manifold structures of  $T_b$  and  $T_v$  proved in [26].

The main results in this paper are Theorem 4 and its corollaries. Having the projection from  $T_b$  onto  $T_v$  compatible with their complex structures, we will prove that the subspace  $T_c$  of chord-arc curves is composed as a union of the fibers of this projection. This can be shown by investigating the set  $T_c$  in view of the group structures of  $T_v$  and  $T_b$ . Then, we see that  $T_c$  and its quotient have desirable structures in this fiber space. The precise statements are put in Section 4, where a motivation of our study of  $T_c$  and related problems are also mentioned.

We organize this paper by dividing it into sections. Some background on the original theory and preliminaries for some necessary material are summarized in Sections 2 and 3, respectively. The new results are all gathered in Section 4. Instead of giving an introduction of these results in the first section, we ask the reader to refer to the content developed in Sections 2 and 3. The rest of the paper after Section 4 is all devoted to the proofs of the main theorems.

## 2. Background on the universal Teichmüller space

In this section, we review the theory of the universal Teichmüller space (see [1, 16, 20] for the details). In particular, we explain the construction and the structure of the quotient space by the little universal Teichmüller space, called the asymptotic Teichmüller space (see [10, 12] for the details). The purpose of this paper is to give an analogue of the asymptotic Teichmüller space for the BMO and VMO Teichmüller spaces defined in the next section.

**2.1. The universal Teichmüller space.** A sense-preserving homeomorphism  $h$  of the unit circle  $\mathbb{S} = \{z \in \mathbb{C} \mid |z| = 1\}$  is said to be *quasisymmetric* if there exists a least positive constant  $C(h)$ , called the quasisymmetry constant of  $h$ , such that the quasisymmetry quotient

$$m_h(x, t) = \frac{|h(e^{i(x+t)}) - h(e^{ix})|}{|h(e^{ix}) - h(e^{i(x-t)})|}$$

takes values in the interval  $[1/C(h), C(h)]$  for all  $x \in [0, 2\pi)$  and  $t \in (0, \pi)$ . Let QS denote the group of all quasisymmetric homeomorphisms of  $\mathbb{S}$ . Beurling and Ahlfors [4] proved that a sense-preserving homeomorphism  $h$  of  $\mathbb{S}$  is quasisymmetric if and only if there exists some quasiconformal homeomorphism of the unit disk  $\mathbb{D} = \{z \in \mathbb{C} \mid |z| < 1\}$  onto itself that has boundary value  $h$ . Later, Douady and Earle [9] gave a quasiconformal extension of a quasisymmetric homeomorphism of  $\mathbb{S}$ , called the Douady–Earle extension, in a conformally natural way.

The *universal Teichmüller space*  $T$  can be defined as the group QS modulo the left action of the group  $\text{Möb}(\mathbb{S})$  of all Möbius transformations of  $\mathbb{S}$ , i.e.,  $T = \text{Möb}(\mathbb{S}) \backslash \text{QS}$ . Alternatively,  $T$  can be identified with the set of representatives  $h \in \text{QS}$  fixing three points, 1,  $-1$  and  $i$ , which can be realized as a subgroup of QS. The topology on  $T$  is induced from that on QS defined by the quasisymmetry constant. This is the real model of the universal Teichmüller space  $T$ .

The universal Teichmüller space  $T$  is also represented by quasiconformal maps. Let

$$M(\mathbb{D}) = \{\mu \in L^\infty(\mathbb{D}) \mid \|\mu\|_\infty < 1\}$$

denote the open unit ball of the Banach space  $L^\infty(\mathbb{D})$  of all essentially bounded measurable functions on  $\mathbb{D}$ . An element in  $M(\mathbb{D})$  is called a *Beltrami coefficient*. By the measurable Riemann mapping theorem, for any  $\mu \in M(\mathbb{D})$ , there is a unique normalized quasiconformal map  $f^\mu$  of  $\mathbb{D}$  onto itself whose complex dilatation is  $\mu$ , where the normalization is given by fixing  $1$ ,  $-1$  and  $i$ . Two elements  $\mu$  and  $\nu$  in  $M(\mathbb{D})$  are equivalent, denoted by  $\mu \sim \nu$ , if  $f^\mu|_{\mathbb{S}} = f^\nu|_{\mathbb{S}}$ . Then, the set  $M(\mathbb{D})/\sim$  of all equivalence classes  $[\mu]$  is the Beltrami coefficient model of  $T$ . Let  $\pi$  be the *Teichmüller projection* from  $M(\mathbb{D})$  onto  $T$  defined by  $\pi(\mu) = [\mu]$ . This is continuous and open, which induces a homeomorphism between  $M(\mathbb{D})/\sim$  and  $\text{Möb}(\mathbb{S}) \setminus \text{QS}$ .

There is also a unique normalized quasiconformal map  $f_\mu$  of the Riemann sphere  $\widehat{\mathbb{C}}$  with complex dilatation  $\mu$  in  $\mathbb{D}$  and  $0$  in  $\mathbb{D}^* = \widehat{\mathbb{C}} - \overline{\mathbb{D}}$ . The normalization of a holomorphic map  $f: \mathbb{D}^* \rightarrow \widehat{\mathbb{C}}$  is given by

$$(1) \quad f(z) = z + \frac{b_1}{z} + \cdots \quad (z \rightarrow \infty).$$

We consider the Bers map  $\Phi: M(\mathbb{D}) \rightarrow B(\mathbb{D}^*)$  sending  $\mu$  to the Schwarzian derivative  $\mathcal{S}(f_\mu|_{\mathbb{D}^*})$  of the conformal map  $f_\mu|_{\mathbb{D}^*}$ , where

$$B(\mathbb{D}^*) = \{\varphi \mid \|\varphi\|_B = \sup_{z \in \mathbb{D}^*} (|z|^2 - 1)^2 |\varphi(z)| < \infty\}$$

is the Banach space of all holomorphic mappings  $\varphi$  on  $\mathbb{D}^* = \widehat{\mathbb{C}} - \overline{\mathbb{D}}$  with the norm  $\|\varphi\|_B$ . The *Bers embedding*  $\beta: T \rightarrow B(\mathbb{D}^*)$  is given by the factorization of the map  $\Phi$  by the Teichmüller projection  $\pi$ , i.e.,  $\beta \circ \pi = \Phi$ . This is a well-defined injection due to the fact that  $f_\mu|_{\mathbb{D}^*} = f_\nu|_{\mathbb{D}^*}$  is equivalent to  $f^\mu|_{\mathbb{S}} = f^\nu|_{\mathbb{S}}$ . It can be proved that  $\Phi$  is a holomorphic split submersion (see [16, Theorem V.5.3], [20, Sections 3.4, 3.5]). In particular, there is a local holomorphic right inverse of  $\Phi$  at every point of  $\Phi(M(\mathbb{D}))$ . This implies that the Bers embedding  $\beta$  is a homeomorphism onto its image, and thus it induces a complex structure of  $T$  as a domain in the Banach space  $B(\mathbb{D}^*)$ . This is the unique complex structure on  $T$  such that the projection  $\pi$  is holomorphic.

It is well known that a quasiconformal homeomorphism of  $\mathbb{D}$  onto itself induces a biholomorphic automorphism of the universal Teichmüller space  $T$ . Precisely, the normalized quasiconformal homeomorphism  $f^\mu$  for  $\mu \in M(\mathbb{D})$  induces a biholomorphic automorphism  $r_\mu: M(\mathbb{D}) \rightarrow M(\mathbb{D})$  which sends  $\nu$  to

$$\nu * \mu^{-1} = \left( \frac{\nu - \mu}{1 - \bar{\mu}\nu} \frac{\partial f^\mu}{\partial \bar{f}^\mu} \right) \circ (f^\mu)^{-1}.$$

This is the complex dilatation of the composition  $f^\nu \circ (f^\mu)^{-1}$ . We denote by  $\nu * \mu$  the complex dilatation of the composition  $f^\nu \circ f^\mu$ , and by  $\mu^{-1}$  the complex dilatation of the inverse  $(f^\mu)^{-1}$ . The map  $r_\mu$  descends down to a biholomorphic automorphism  $R_{[\mu]}$  of  $T$  defined by  $R_{[\mu]} \circ \pi = \pi \circ r_\mu$  (see [16, Section V.5.4], [20, Section 3.6.2]).

The group operation  $*$  on  $M(\mathbb{D})$  also descends down to the operation  $*$  on  $T$  by  $[\mu] * [\nu] = [\mu * \nu]$ . Combined with the inverse operation  $[\mu]^{-1} = [\mu^{-1}]$ , this turns out to be the group structure  $(T, *)$  of the universal Teichmüller space, which is the same as the group structure of  $T$  regarded as a subgroup of QS. Then,  $R_{[\mu]}$  is the right translation of  $T$  by  $[\mu]$  which sends  $[\nu]$  to  $[\nu] * [\mu]^{-1}$ .

**2.2. The asymptotic Teichmüller space.** A quasymmetric homeomorphism  $h \in \text{QS}$  of  $\mathbb{S}$  is called *symmetric* if, in addition, the quasimetric quotient satisfies  $m_h(x, t) \rightarrow 1$  as  $t \rightarrow 0$  uniformly. We denote the subgroup of QS consisting of all symmetric homeomorphisms of  $\mathbb{S}$  by  $\text{Sym}$ . It was proved by Gardiner and Sullivan [13] that  $\text{Sym}$  is the characteristic topological subgroup of QS. This in particular implies that, for any  $h \in \text{Sym}$ ,  $f_n$  converges to  $f$  in QS if and only if  $h \circ f_n$  converges to  $h \circ f$ . The *little universal Teichmüller space* is defined by  $T_0 = \text{Möb}(\mathbb{S}) \setminus \text{Sym}$ . This can be regarded as a closed topological subgroup of  $(T, *)$ .

It is known that  $h$  is symmetric if and only if  $h$  can be extended to an asymptotically conformal homeomorphism  $f$  of the unit disk  $\mathbb{D}$  onto itself. In fact, the Douady–Earle extension of  $h$  is asymptotically conformal when  $h$  is symmetric. Here, by an *asymptotically conformal homeomorphism*  $f$  of the unit disk  $\mathbb{D}$ , we mean that its complex dilatation  $\mu$  vanishes at the boundary, that is,  $\text{ess. sup}_{r \leq |z| < 1} |\mu(z)| \rightarrow 0$  as  $r \rightarrow 1$ . The closed subspace of  $M(\mathbb{D})$  consisting of all Beltrami coefficients vanishing at the boundary is denoted by  $M_0(\mathbb{D})$ . Then,  $\pi(M_0(\mathbb{D}))$  is a closed subspace of  $T$ , which coincides with  $T_0$ . The image of  $M_0(\mathbb{D})$  under the Bers map  $\Phi: M(\mathbb{D}) \rightarrow B(\mathbb{D})$  is contained in the Banach subspace  $B_0(\mathbb{D}^*)$  of  $B(\mathbb{D}^*)$  consisting of those holomorphic functions  $\varphi$  such that  $(|z|^2 - 1)^2 |\varphi(z)| \rightarrow 0$  as  $|z| \rightarrow 1$  uniformly. Moreover, the local holomorphic inverse of  $\Phi$  at every point of  $\Phi(M_0(\mathbb{D}))$  maps its neighborhood in  $B_0(\mathbb{D}^*)$  to  $M_0(\mathbb{D})$ . Thus,  $T_0$  has the structure of the complex Banach submanifold of  $T$  modeled on  $B_0(\mathbb{D}^*)$ . In particular, we have that  $\beta(T_0) = \Phi(M_0(\mathbb{D})) = \beta(T) \cap B_0(\mathbb{D}^*)$ .

We consider the quotient space  $AT = T_0 \setminus T$ , which is called the *asymptotic Teichmüller space* (see [10, 13]). If we regard  $T = \text{Möb}(\mathbb{S}) \setminus \text{QS}$  and  $T_0 = \text{Möb}(\mathbb{S}) \setminus \text{Sym}$ , then  $AT$  can be realized as the set of all cosets  $\text{Sym} \setminus \text{QS}$  in the group QS. Alternatively, the equivalence class in  $T$  under the quotient of  $T_0$  containing  $\tau \in T$  is given by  $R_\tau^{-1}(T_0)$ . In order to introduce a complex structure to  $AT$  modeled on the quotient Banach space  $B_0(\mathbb{D}^*) \setminus B(\mathbb{D}^*)$ , we have to verify the compatibility of the quotients  $T_0 \setminus T$  and  $B_0(\mathbb{D}^*) \setminus B(\mathbb{D}^*)$  under the Bers embedding  $\beta: T \rightarrow B(\mathbb{D}^*)$ .

It was proved in [13] that the quotient Bers embedding

$$\widehat{\beta}: T_0 \setminus T \rightarrow B_0(\mathbb{D}^*) \setminus B(\mathbb{D}^*)$$

is well-defined and locally injective. Then,  $AT = T_0 \setminus T$  becomes a complex manifold having local coordinates in the quotient Banach space  $B_0(\mathbb{D}^*) \setminus B(\mathbb{D}^*)$ . Later, Kahn (see [12, Section 16.8]) proved further that  $\widehat{\beta}$  is globally injective. Earle, Markovic and Saric [11, Theorem 4] gave a different proof for this by using the Douady–Earle extension and generalized the result in a compatible way with Fuchsian group action. As in the following theorem, we can formulate these claims so that the decomposition of  $T$  into the submanifolds  $R_\tau^{-1}(T_0)$  corresponds bijectively to the decomposition of  $\beta(T) \subset B(\mathbb{D}^*)$  into the intersections of the affine subspaces isometric to  $B_0(\mathbb{D}^*)$ . We call this decomposition the *affine foliated structure* of  $T$  induced by  $T_0$ . See [18] for a comprehensive exposition.

**Theorem A.**  $\beta \circ R_\tau^{-1}(T_0) = \beta(T) \cap \{B_0(\mathbb{D}^*) + \beta(\tau)\}$  for every  $\tau \in T$ .

The inclusion  $\subset$  in the above equality implies that the quotient Bers embedding  $\widehat{\beta}$  is well-defined. The converse inclusion  $\supset$  implies that  $\widehat{\beta}$  is injective. Combining the well-definedness and the injectivity of  $\widehat{\beta}$  with the homeomorphy of  $\beta$  from  $T$  onto its image in  $B(\mathbb{D}^*)$ , we have the following result naturally.

**Corollary B.** *The quotient Bers embedding  $\widehat{\beta}: T_0 \setminus T \rightarrow B_0(\mathbb{D}^*) \setminus B(\mathbb{D}^*)$  is a homeomorphism onto its image. Consequently,  $AT = T_0 \setminus T$  possesses a complex*

structure such that  $\hat{\beta}$  is a biholomorphic homeomorphism from  $T_0 \setminus T$  onto its image in  $B_0(\mathbb{D}^*) \setminus B(\mathbb{D}^*)$ .

In this paper, we will obtain the analogous results to these two claims in the setting of the BMO theory of the universal Teichmüller space.

### 3. Preliminaries on the BMO and VMO Teichmüller spaces

In this section, we give basic definitions and fundamental results on the BMO theory of the universal Teichmüller space (see [3, 25, 26] for the details). We explain these in an analogous way to those for the universal Teichmüller space given in the previous section.

**3.1. The BMO Teichmüller space.** A sense-preserving homeomorphism  $h$  of the unit circle  $\mathbb{S}$  is called *strongly quasisymmetric* if for any  $\varepsilon > 0$  there is some  $\delta > 0$  such that for any arc  $I \subset \mathbb{S}$  and any Borel set  $E \subset I$ ,

$$|E| \leq \delta |I| \Rightarrow |h(E)| \leq \varepsilon |h(I)|.$$

By this definition, a strongly quasisymmetric homeomorphism is quasisymmetric, and the composition of strongly quasisymmetric homeomorphisms is strongly quasisymmetric. Moreover, it is known that the inverse of a strongly quasisymmetric homeomorphism is strongly quasisymmetric (see [7]). We denote by SQS the subgroup of QS consisting of all strongly quasisymmetric homeomorphisms of  $\mathbb{S}$ .

Any strongly quasisymmetric homeomorphism  $h$  is absolutely continuous with  $\log h' \in \text{BMO}(\mathbb{S})$  (see [14, Chap. 6]), but the converse is not true. Here, an integrable function  $\phi$  on  $\mathbb{S}$  belongs to  $\text{BMO}(\mathbb{S})$  if

$$\|\phi\|_{\text{BMO}} = \sup_{I \subset \mathbb{S}} \frac{1}{|I|} \int_I |\phi(e^{i\theta}) - \phi_I| \frac{d\theta}{2\pi} < \infty,$$

where the supremum is taken over all arcs  $I$  on  $\mathbb{S}$ ,  $|I| = \int_I d\theta/2\pi$  is the normalized length of  $I$ , and  $\phi_I$  denotes the average of  $\phi$  over  $I$ . The *BMO Teichmüller space* is defined by  $T_b = \text{Möb}(\mathbb{S}) \setminus \text{SQS}$ . This can be regarded as a subgroup of  $(T, *)$ . The topology on  $T_b$  is induced from that on SQS given by the BMO norm, that is, by the distance  $d(h_1, h_2) = \|\log h'_1 - \log h'_2\|_{\text{BMO}}$ .

As in the case of the universal Teichmüller space, the BMO Teichmüller space  $T_b$  has the corresponding space for Beltrami coefficients. A measure  $\lambda = \lambda(z) dx dy$  on  $\mathbb{D}$  is called a *Carleson measure* if

$$\|\lambda\|_c = \sup \frac{\lambda(S_{h,\theta_0})}{h} < \infty$$

where the supremum is taken over all sectors

$$S_{h,\theta_0} = \{re^{i\theta} \in \mathbb{D} \mid 1-h \leq r < 1, |\theta - \theta_0| \leq \pi h\}$$

for  $h \in (0, 1]$  and  $\theta_0 \in [0, 2\pi)$ . We denote by  $\text{CM}(\mathbb{D})$  the set of all Carleson measures on  $\mathbb{D}$ . For  $\mathbb{D}^* = \widehat{\mathbb{C}} - \mathbb{D}$ , the set  $\text{CM}(\mathbb{D}^*)$  of the Carleson measures on  $\mathbb{D}^*$  can be defined similarly. For  $\mu \in L^\infty(\mathbb{D})$  and for the Poincaré density  $\rho_{\mathbb{D}}(z) = (1 - |z|^2)^{-1}$  (with curvature constant equal to  $-4$ ) on  $\mathbb{D}$ , we set

$$\lambda_\mu = |\mu(z)|^2 \rho_{\mathbb{D}}(z) dx dy.$$

Then the linear subspace  $\mathcal{L}(\mathbb{D}) \subset L^\infty(\mathbb{D})$  consisting of all  $\mu$  with  $\lambda_\mu \in \text{CM}(\mathbb{D})$  is a Banach space with a norm  $\|\mu\|_* = \|\mu\|_\infty + \|\lambda_\mu\|_c^{1/2}$ . Moreover, we consider the corresponding space of Beltrami coefficients as  $\mathcal{M}(\mathbb{D}) = M(\mathbb{D}) \cap \mathcal{L}(\mathbb{D})$ . Then,  $T_b$  is the image of  $\mathcal{M}(\mathbb{D})$  under the Teichmüller projection  $\pi: M(\mathbb{D}) \rightarrow T$ . The quotient

topology on  $T_b$  induced from  $\mathcal{M}(\mathbb{D})$  by  $\pi$  coincides with the topology on  $T_b$  induced from the BMO norm.

There is also a subspace of holomorphic mappings corresponding to  $T_b$ . For  $\varphi \in B(\mathbb{D}^*)$ , another norm is given by  $\|\varphi\|_{\mathcal{B}} = \|\tilde{\lambda}_\varphi\|_c^{1/2}$ , where

$$\tilde{\lambda}_\varphi = |\varphi(z)|^2 \rho_{\mathbb{D}^*}^{-3}(z) dx dy$$

is a Carleson measure on  $\mathbb{D}^*$  for the Poincaré density  $\rho_{\mathbb{D}^*}(z) = (|z|^2 - 1)^{-1}$ . We consider the Banach space  $\mathcal{B}(\mathbb{D}^*) \subset B(\mathbb{D}^*)$  consisting of all such elements  $\varphi$  that  $\tilde{\lambda}_\varphi \in \text{CM}(\mathbb{D}^*)$  equipped with the norm  $\|\varphi\|_{\mathcal{B}}$ . The following result was proved in [26, Theorem 5.1].

**Theorem C.** *The Bers map  $\Phi$  restricted to  $\mathcal{M}(\mathbb{D})$  is a holomorphic map into  $\mathcal{B}(\mathbb{D}^*)$  with a local holomorphic right inverse at every point of the image  $\Phi(\mathcal{M}(\mathbb{D}))$ . The Bers embedding  $\beta$  of  $T_b$  is a homeomorphism onto the domain  $\beta(T_b) = \Phi(\mathcal{M}(\mathbb{D}))$  in  $\mathcal{B}(\mathbb{D}^*)$ . In particular,  $T_b$  has a complex structure modeled on  $\mathcal{B}(\mathbb{D}^*)$ .*

**3.2. The VMO Teichmüller space.** We say that a strongly quasymmetric homeomorphism  $h \in \text{SQS}$  is *strongly symmetric* if  $\log h' \in \text{VMO}(\mathbb{S})$ , where a function  $\phi \in \text{BMO}(\mathbb{S})$  belongs to  $\text{VMO}(\mathbb{S})$  if

$$\lim_{|I| \rightarrow 0} \frac{1}{|I|} \int_I |\phi(e^{i\theta}) - \phi_I| \frac{d\theta}{2\pi} = 0$$

uniformly. In fact,  $\text{VMO}(\mathbb{S})$  is a closed subspace of  $\text{BMO}(\mathbb{S})$ , and it is precisely the closure of the space of all continuous functions on  $\mathbb{S}$  under the BMO topology (see [23]). We denote by  $\text{SS}$  the subgroup of  $\text{SQS}$  consisting of all strongly symmetric homeomorphisms. It is easy to see that the inclusion relation  $\text{SS} \subset \text{Sym}$  holds. It was proved in [28] that  $\text{SS}$  is the characteristic topological subgroup of  $\text{SQS}$ . This in particular implies that, for any  $h \in \text{SS}$ ,  $f_n$  converges to  $f$  in  $\text{SQS}$  if and only if  $h \circ f_n$  converges to  $h \circ f$ . The *VMO Teichmüller space*  $T_v$  is defined to be  $T_v = \text{Möb}(\mathbb{S}) \setminus \text{SS}$ , which can be regarded as a closed topological subgroup of  $(T_b, *)$ .

A Carleson measure  $\lambda \in \text{CM}(\mathbb{D})$  is called a *vanishing Carleson measure* if

$$\lim_{h \rightarrow 0} \frac{\lambda(S_{h, \theta_0})}{h} = 0$$

uniformly for  $\theta_0 \in [0, 2\pi)$ . We denote the set of all vanishing Carleson measures on  $\mathbb{D}$  by  $\text{CM}_0(\mathbb{D})$ . The set  $\text{CM}_0(\mathbb{D}^*)$  of the vanishing Carleson measures on  $\mathbb{D}^*$  can be defined similarly. Let  $\mathcal{M}_0(\mathbb{D})$  be the closed subspace of  $\mathcal{M}(\mathbb{D})$  consisting of all Beltrami coefficients  $\mu$  such that  $\lambda_\mu \in \text{CM}_0(\mathbb{D})$ . Then,  $\pi(\mathcal{M}_0(\mathbb{D}))$  is a closed subspace of  $T_b$ , which coincides with  $T_v$ . We denote by  $\mathcal{B}_0(\mathbb{D}^*)$  the Banach subspace of  $\mathcal{B}(\mathbb{D}^*)$  consisting of all elements  $\varphi$  such that  $\tilde{\lambda}_\varphi = |\varphi(z)|^2 \rho_{\mathbb{D}^*}^{-3}(z) dx dy \in \text{CM}_0(\mathbb{D}^*)$ . Then  $\mathcal{B}_0(\mathbb{D}^*) \subset \mathcal{B}(\mathbb{D}^*)$  by [26, Lemma 4.1]. It was proved in [26, Theorems 4.1, 5.2] that  $\Phi$  maps  $\mathcal{M}_0(\mathbb{D})$  into  $\mathcal{B}_0(\mathbb{D}^*)$  and the Bers embedding  $\beta$  of  $T_v$  is a homeomorphism onto a domain  $\beta(T_v) = \Phi(\mathcal{M}_0(\mathbb{D}))$  in  $\mathcal{B}_0(\mathbb{D}^*)$ .

**3.3. The chord-arc curve space.** A rectifiable closed Jordan curve  $\Gamma$  in the complex plane  $\mathbb{C}$  is called a *chord-arc curve* if

$$(2) \quad l_\Gamma(z_1, z_2) / |z_1 - z_2| \leq K$$

for any  $z_1, z_2 \in \Gamma$ , where  $l_\Gamma(z_1, z_2)$  denotes the Euclidean length of the shorter arc of  $\Gamma$  between  $z_1$  and  $z_2$ . The smallest such constant  $K > 1$  is called the chord-arc constant for  $\Gamma$ . A chord-arc curve is in particular a quasicircle. Further, if the ratio

in (2) tends to 1 uniformly as  $|z_1 - z_2| \rightarrow 0$ , then the curve  $\Gamma$  is called *asymptotically smooth* in the sense of Pommerenke [21].

The following fact is well-known as the characterization of a chord-arc curve (see [15, Proposition 1.13]).

**Proposition D.** *A chord-arc curve is the image of  $\mathbb{S}$  under a bi-Lipschitz homeomorphism  $f$  of  $\mathbb{C}$  onto itself with respect to the Euclidean distance. That is, there exists a homeomorphism  $f: \mathbb{C} \rightarrow \mathbb{C}$  with a constant  $C \geq 1$  such that  $f(\mathbb{S}) = \Gamma$  and  $C^{-1}|z - w| \leq |f(z) - f(w)| \leq C|z - w|$  for all  $z, w \in \mathbb{C}$ .*

If  $\Gamma$  is a simple curve passing through  $\infty$  satisfying the chord-arc condition (2), we can transfer it to a bounded chord-arc curve by a Möbius transformation because the chord-arc condition is Möbius invariant, or equivalently, the distance in the chord-arc condition can be replaced with the spherical distance (see [17, p. 877]). This also implies that Proposition D can be stated equivalently for an unbounded chord-arc curve by the same bi-Lipschitz condition.

Let  $\Gamma$  be a bounded Jordan curve in the Riemann sphere  $\widehat{\mathbb{C}}$ , let  $\Omega$  and  $\Omega^*$  denote its inner and outer domains in  $\widehat{\mathbb{C}}$ , respectively, and let  $g$  and  $f$  be conformal maps of  $\mathbb{D}$  and  $\mathbb{D}^*$  onto  $\Omega$  and  $\Omega^*$ , respectively. We always assume that  $f: \mathbb{D}^* \rightarrow \Omega^*$  is normalized so that it satisfies (1). These two maps extend homeomorphically to the boundary, and hence  $h = (g|_{\mathbb{S}})^{-1} \circ (f|_{\mathbb{S}})$  determines a sense-preserving homeomorphism of  $\mathbb{S}$  onto itself, which is called the *conformal welding homeomorphism* with respect to  $\Gamma$ .

It is well known that  $h$  is strongly quasimetric if and only if the curve  $\Gamma$  is a quasicircle satisfying the so-called Bishop–Jones condition (see [6]): for any  $z \in \Omega$  there exists a domain  $\Omega_z (\subset \Omega)$  containing  $z$  bounded by a chord-arc curve with constant  $K$  such that the diameter of  $\Omega_z$  is uniformly comparable to  $\text{dist}(z, \Gamma)$  and the Hausdorff linear measure of  $\Gamma \cap \partial\Omega_z$  is bounded from below by  $C \text{dist}(z, \Gamma)$ , where  $K > 1$  and  $C > 0$  depend only on  $\Gamma$ . The Bishop–Jones condition is invariant under a bi-Lipschitz homeomorphism of  $\mathbb{C}$  onto itself in the Euclidean metric, and hence, any chord-arc curve satisfies this condition. It was proved in [21] that  $h$  is strongly symmetric if and only if the curve  $\Gamma$  is asymptotically smooth. However, although chord-arc curves are in a very special class of quasicircles, no characterization has been found in terms of their conformal welding homeomorphisms of  $\mathbb{S}$ .

We denote the set of all these conformal welding homeomorphisms for chord-arc curves by CQS. Then, we have the following proper inclusion relations:  $\text{SS} \subsetneq \text{CQS} \subsetneq \text{SQS}$ . Here, the strictness of the second inclusion is seen by a fact that a quasicircle satisfying the Bishop–Jones condition is not necessarily rectifiable (see [5, 24]). Similarly to  $T_b = \text{Möb}(\mathbb{S}) \setminus \text{SQS}$  and  $T_v = \text{Möb}(\mathbb{S}) \setminus \text{SS}$ , we define  $T_c = \text{Möb}(\mathbb{S}) \setminus \text{CQS}$ . The following fact was essentially shown by Zinsmeister [31] (see also [3]).

**Proposition E.**  *$T_c$  is an open subset of  $T_b$  containing  $T_v$ .*

#### 4. Statement of the results

We state our results in this paper, which fall into two parts. The first part is concerning the affine foliated structure of the BMO Teichmüller space  $T_b$  by the VMO Teichmüller space  $T_v$ . This is an analogous result with Theorem A, which gives a foundation to investigate the structure of the quotient Teichmüller space.

**Theorem 1.**  *$\beta \circ R_\tau^{-1}(T_v) = \beta(T_b) \cap \{\mathcal{B}_0(\mathbb{D}^*) + \beta(\tau)\}$  for every  $\tau \in T_b$ .*

We note that  $R_\tau^{-1}(T_v)$  for each  $\tau \in T_b$  is an equivalence class of the quotient space  $T_v \setminus T_b \cong \text{SS} \setminus \text{SQS}$  containing  $\tau$ , which is also a closed subspace of  $T_b$  biholomorphically equivalent to  $T_v$ . By this theorem, we have the decomposition of the Bers embedding as

$$\beta(T_b) = \bigsqcup_{[\tau] \in T_v \setminus T_b} \beta \circ R_\tau^{-1}(T_v) = \bigsqcup_{[\psi] \in \mathcal{B}_0(\mathbb{D}^*) \setminus \mathcal{B}(\mathbb{D}^*)} \beta(T_b) \cap \{\mathcal{B}_0(\mathbb{D}^*) + \psi\}.$$

Based on Theorems 1 and C, the quotient space  $T_v \setminus T_b$  is provided with a complex structure modeled on the quotient Banach space  $\mathcal{B}_0(\mathbb{D}^*) \setminus \mathcal{B}(\mathbb{D}^*)$ . The argument for this is the same as that for Corollary B in the case of the asymptotic Teichmüller space.

**Corollary 2.** *The quotient Bers embedding*

$$\widehat{\beta}: T_v \setminus T_b \rightarrow \mathcal{B}_0(\mathbb{D}^*) \setminus \mathcal{B}(\mathbb{D}^*)$$

is well-defined and injective to be a homeomorphism of  $T_v \setminus T_b$  onto its image. Consequently,  $T_v \setminus T_b$  possesses a complex structure such that  $\widehat{\beta}$  is biholomorphic.

Moreover, we have the following result on biholomorphic automorphisms of  $T_v \setminus T_b$  with respect to its complex structure. This is well-known in the theory of asymptotic Teichmüller space (see [19, Proposition 4.1]).

**Corollary 3.** *Let  $p: T_b \rightarrow T_v \setminus T_b$  be the quotient projection from  $T_b$  onto  $T_v \setminus T_b$ . For every  $\tau \in T_b$ , the biholomorphic automorphism  $R_\tau$  of  $T_b$  induces a biholomorphic automorphism  $\widehat{R}_\tau$  of  $T_v \setminus T_b$  satisfying  $p \circ R_\tau = \widehat{R}_\tau \circ p$ .*

The second part of our results is concerning the structure of the space of chord-arc curves. We will show that CQS does not carry a group structure under the composition. This follows from the claim that every element of SQS can be represented as a finite composition of elements in CQS. However, CQS is preserved under the inverse operation and under the left and right actions of SS. In particular, CQS is preserved under the conjugation by SS. We state these claims in the framework of Teichmüller spaces; the chord-arc curve space is identified with a subspace  $T_c = \text{Möb}(\mathbb{S}) \setminus \text{CQS}$  of the BMO Teichmüller space  $(T_b, *)$ . For each  $\sigma \in T_b$ , the left translation  $L_\sigma: T_b \rightarrow T_b$  is defined by  $L_\sigma(\tau) = \sigma * \tau$  for every  $\tau \in T_b$ .

**Theorem 4.** *The following statements hold.*

- (a) *Each element of  $T_b$  can be represented as a finite composition of elements in  $T_c$ . Hence,  $T_c$  is not a subgroup of  $T_b$ .*
- (b) *The inverse element  $\tau^{-1}$  belongs to  $T_c$  for every  $\tau \in T_c$ .*
- (c)  *$L_\sigma(T_c) = T_c$  and  $R_\sigma(T_c) = T_c$  for every  $\sigma \in T_v$ .*

**Remark.** Statement (a) is a general fact not only for  $U = T_c$  but also for any open neighborhood  $U \subsetneq T_b$  of the origin  $o = [0]$  of the Teichmüller space. This is seen from its proof. Statement (b) does not assert that the inverse operation  $\tau \mapsto \tau^{-1}$  is continuous. In fact, this is not necessarily continuous on  $T_c$  but continuous on  $T_v$ . For statement (c), we have only to show the inclusion  $L_\sigma(T_c) \subset T_c$ . The inverse inclusion follows from the facts that  $L_\sigma^{-1} = L_{\sigma^{-1}}$  and  $\sigma^{-1} \in T_v$ . The equality for the right translation is obtained by taking the inverse of the equality for the left translation and applying statement (b).

The above condition  $L_\sigma(T_c) \subset T_c$  for every  $\sigma \in T_v$  is equivalent to that  $R_\tau^{-1}(T_v) \subset T_c$  for every  $\tau \in T_c$ . We verify this property in the proof. Concerning the fiber structure of  $T_c$  with respect to the projection  $p: T_b \rightarrow T_v \setminus T_b$ , this condition implies



that  $T_c$  consists of all fibers of  $p$  that intersect  $T_c$ . Moreover, if we consider this as the property of the right translation  $R_\sigma$  by  $\sigma \in T_v$  preserving  $T_c$ , the projection  $p$  restricted to  $T_c$  is also obtained as the quotient by the group action of  $T_v$ .

**Corollary 5.** *For each  $\tau \in T_c$ , the fiber  $R_\tau^{-1}(T_v)$  of the projection  $p: T_b \rightarrow T_v \setminus T_b$  is entirely contained in  $T_c$ . Hence,  $T_c = \bigsqcup_{[\tau] \in p(T_c)} R_\tau^{-1}(T_v)$ . For each  $\sigma \in T_v$ , the right translation  $R_\sigma$  acts on  $T_c$  as a biholomorphic automorphism, and  $p(T_c)$  is given as the quotient  $T_v \setminus T_c$  of this group action.*

**Problem.** We have seen that  $T_v$  acts on  $T_c$  as a group of biholomorphic automorphisms. Then, we may ask about whether this property characterizes  $T_v$ , namely, if the stabilizer subgroup

$$\{\tau \in T_b \mid R_\tau(T_c) = T_c = L_\tau(T_c)\}$$

containing  $T_v$  coincides with  $T_v$  or not. As a related question, for any  $\tau \in T_c - T_v$ , we ask about the existence of an integer  $n$  such that  $\tau^n = \tau * \dots * \tau \notin T_c$ .

Corollary 5 implies the following equation in the Bers embedding. Combined with Theorem 1, this yields the affine foliated structure of  $T_c$  by  $T_v$ .

**Corollary 6.**  $\beta(T_b) \cap \{\mathcal{B}_0(\mathbb{D}^*) + \beta(\tau)\} = \beta(T_c) \cap \{\mathcal{B}_0(\mathbb{D}^*) + \beta(\tau)\}$  for every  $\tau \in T_c$ .

The quotient Bers embedding from  $T_v \setminus T_c$  into  $\mathcal{B}_0(\mathbb{D}^*) \setminus \mathcal{B}(\mathbb{D}^*)$ , considered in [29, Theorem 2.2], is well-defined and injective. This can be extended to the embedding of  $T_v \setminus T_b$  as in Corollary 2. Then, we provide the quotient space  $T_v \setminus T_c$  with a complex structure modeled on the quotient Banach space  $\mathcal{B}_0(\mathbb{D}^*) \setminus \mathcal{B}(\mathbb{D}^*)$ .

**Corollary 7.** *The quotient Bers embedding  $\widehat{\beta}$  maps  $T_v \setminus T_c$  homeomorphically onto its image in  $\mathcal{B}_0(\mathbb{D}^*) \setminus \mathcal{B}(\mathbb{D}^*)$ , and  $T_v \setminus T_c$  possesses a complex structure as a domain in  $T_v \setminus T_b$ .*

Connectivity of  $T_c$  is an open problem (see [2], [3, p. 614]). Since  $T_v$  is connected, Corollary 5 also implies the following reduction on this problem.

**Corollary 8.**  $T_c$  is connected if and only if  $T_v \setminus T_c$  is connected.

Corollaries 5, 6, 7, and 8 follow directly from the preceding results. The remainder of this paper is devoted to the proofs of Theorem 1 with Corollaries 2 and 3 and Theorem 4.

## 5. Proof of Theorem 1

For every  $\tau \in T_b$ , let  $f^\nu: \mathbb{D} \rightarrow \mathbb{D}$  be a normalized quasiconformal extension of  $\tau$  with complex dilatation  $\nu \in \mathcal{M}(\mathbb{D})$  (i.e.  $\pi(\nu) = \tau$ ) that is bi-Lipschitz under the Poincaré metric on  $\mathbb{D}$  (for instance, the Douady–Earle extension of  $\tau$  satisfies this condition; see [8]). Let  $\psi = \Phi(\nu) \in \mathcal{B}(\mathbb{D}^*)$ .

**5.1. Proof of the inclusion  $\subset$ .** We divide the arguments into two steps. We first deal with the special case that  $\mu \in \mathcal{M}_0(\mathbb{D})$  has a compact support. Then, we extend this to the general case by means of an approximation process.

We take such a Beltrami coefficient  $\mu \in \mathcal{M}_0(\mathbb{D})$  with compact support. We will show that  $\Phi(\mu * \nu) - \Phi(\nu) \in \mathcal{B}_0(\mathbb{D}^*)$ . Then, the inclusion  $\subset$  follows from

$$\Phi(\mu * \nu) - \Phi(\nu) = \beta \circ \pi(\mu * \nu) - \beta \circ \pi(\nu) = \beta \circ R_\tau^{-1}(\pi(\mu)) - \beta(\tau).$$

Let  $f_{\mu*\nu}: \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$  be the quasiconformal homeomorphism with complex dilatation  $\mu * \nu$  on  $\mathbb{D}$  that is conformal on  $\mathbb{D}^*$ . Set  $\widehat{f} = f_{\mu*\nu} \circ f_\nu^{-1}$ . Then,  $\widehat{f}$  is a quasiconformal

homeomorphism with complex dilatation  $\widehat{\mu}$  on  $\Omega = f_\nu(\mathbb{D})$  whose support is contained in a Jordan domain  $\Omega_0$  with  $\overline{\Omega_0} \subset \Omega$ , and is conformal on  $\Omega^* = f_\nu(\mathbb{D}^*)$  with

$$(3) \quad |\mathcal{S}(\widehat{f})|^2 \rho_{\Omega^*}^{-3} = \left( |\mathcal{S}(f_{\mu*\nu}) - \mathcal{S}(f_\nu)|^2 \rho_{\mathbb{D}^*}^{-3} \right) \circ f_\nu^{-1} |(f_\nu^{-1})'|.$$

In fact,  $\widehat{f}$  is conformal on the larger domain  $\Omega_0^* = \widehat{\mathbb{C}} - \overline{\Omega_0}$ .

It is known that  $|\mathcal{S}(\widehat{f})(z)| \rho_{\Omega_0^*}^{-2}(z) \leq 12$  for  $z \in \Omega_0^*$  (see [16, p. 67]). Since the Poincaré density is monotone with respect to the domain, we have  $\rho_{\Omega_0^*}(z) \leq \rho_{\Omega^*}(z)$ . Then, there exists a constant  $C$  such that

$$|\mathcal{S}(\widehat{f})(z)|^2 \rho_{\Omega^*}^{-3}(z) \leq 144 \rho_{\Omega_0^*}^4(z) \rho_{\Omega^*}^{-3}(z) \leq 144 \rho_{\Omega_0^*}(z) \leq C$$

for  $z \in \Omega^*$ . From this, we deduce that  $|\mathcal{S}(\widehat{f})|^2 \rho_{\Omega^*}^{-3} \in \text{CM}_0(\Omega^*)$ . By (3) and well-definedness of the pull-back operator from  $\text{CM}_0(\Omega^*)$  into  $\text{CM}_0(\mathbb{D}^*)$  (see [29, Theorem 3.1]), we have that  $|\mathcal{S}(f_{\mu*\nu}) - \mathcal{S}(f_\nu)|^2 \rho_{\mathbb{D}^*}^{-3} \in \text{CM}_0(\mathbb{D}^*)$ , and thus  $\Phi(\mu*\nu) - \Phi(\nu) \in \mathcal{B}_0(\mathbb{D}^*)$ .

Next, we consider the general case. For any  $\sigma \in T_\nu$ , the complex dilatation of the Douady–Earle extension of  $\sigma$  is denoted by  $\mu$ . Then,  $\mu \in \mathcal{M}_0(\mathbb{D})$  by [27, Theorem 3.7] (see also [30]). We take an increasing sequence of positive numbers  $r_n < 1$  ( $n = 1, 2, \dots$ ) tending to 1. Let  $\Delta_n$  be an open disk of radius  $r_n$  centered at the origin, and set  $A_n = \mathbb{D} - \overline{\Delta_n}$ . We define

$$\mu_n = \begin{cases} \mu & \text{on } \overline{\Delta_n}, \\ 0 & \text{on } A_n. \end{cases}$$

Then,  $\{\mu_n\}$  is a sequence of complex dilatations with compact support such that

$$(4) \quad \|\mu - \mu_n\|_* = \|\mu - \mu_n\|_\infty + \|\lambda_{\mu - \mu_n}\|_c^{1/2} = \|\mu|_{A_n}\|_\infty + \|\lambda_{\mu|_{A_n}}\|_c^{1/2} \rightarrow 0$$

as  $n \rightarrow \infty$ . Indeed, it was proved in [11] that the complex dilatation of the Douady–Earle extension of a symmetric homeomorphism is in  $M_0(\mathbb{D})$ . Combining this with the inclusion relation  $\text{SS} \subset \text{Sym}$ , we see that  $\mu$  belongs to  $M_0(\mathbb{D})$ , which implies that the first term of the second line of (4) tends to 0. By the definition of  $\mathcal{M}_0(\mathbb{D})$ , the second term also tends to 0.

Since  $f_\nu$  is bi-Lipschitz under the Poincaré metric as we mentioned at the beginning of this section,  $\nu$  induces a biholomorphic automorphism  $r_\nu^{-1}: \mathcal{M}(\mathbb{D}) \rightarrow \mathcal{M}(\mathbb{D})$  (see [26, Remark 5.1]). Then, we have

$$\|r_\nu^{-1}(\mu) - r_\nu^{-1}(\mu_n)\|_* = \|\mu * \nu - \mu_n * \nu\|_* \rightarrow 0$$

as  $n \rightarrow \infty$ . The continuity of  $\Phi$  yields that

$$\|(\Phi(\mu * \nu) - \Phi(\nu)) - (\Phi(\mu_n * \nu) - \Phi(\nu))\|_{\mathcal{B}} = \|\Phi(\mu * \nu) - \Phi(\mu_n * \nu)\|_{\mathcal{B}} \rightarrow 0$$

as  $n \rightarrow \infty$ . We have proved that  $\Phi(\mu_n * \nu) - \Phi(\nu) \in \mathcal{B}_0(\mathbb{D}^*)$  in the first step. Then, it follows that  $\Phi(\mu * \nu) - \Phi(\nu) \in \mathcal{B}_0(\mathbb{D}^*)$  from the fact that  $\mathcal{B}_0(\mathbb{D}^*)$  is closed in  $\mathcal{B}(\mathbb{D}^*)$ . This proves the inclusion  $\subset$ .  $\square$

**5.2. Proof of the inclusion  $\supset$ .** This can be proved by using the following claim, which is shown in [18, Proposition 3.3].

**Claim.** Let  $f_\nu: \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$  be a quasiconformal homeomorphism with complex dilatation  $\nu \in M(\mathbb{D})$  that is bi-Lipschitz between  $\mathbb{D}$  and  $\Omega = f_\nu(\mathbb{D})$  under their Poincaré metrics, and is conformal on  $\mathbb{D}^*$  with  $\mathcal{S}(f_\nu|_{\mathbb{D}^*}) = \psi$ . Then, for every  $\varphi \in B_0(\mathbb{D}^*)$ , there exists a quasiconformal homeomorphism  $\widehat{f}: \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$  with complex dilatation  $\widehat{\mu}$  on  $\Omega$

vanishing at the boundary that is conformal on  $\Omega^* = f_\nu(\mathbb{D}^*)$  with  $\mathcal{S}(\widehat{f} \circ f_\nu|_{\mathbb{D}^*}) = \varphi + \psi$  such that the following statements are valid:  $\widehat{f}$  is decomposed into two quasiconformal homeomorphisms  $\widehat{f}_0$  and  $\widehat{f}_1$  of  $\widehat{\mathbb{C}}$  with  $\widehat{f} = \widehat{f}_0 \circ \widehat{f}_1$ , where  $\widehat{f}_1$  is conformal on  $\Omega^*$  with  $\mathcal{S}(\widehat{f}_1 \circ f_\nu|_{\mathbb{D}^*}) = \varphi_1 + \psi$ , satisfying the following properties:

(i) the complex dilatation  $\widehat{\mu}_1$  of  $\widehat{f}_1$  on  $\Omega$  satisfies

$$|\widehat{\mu}_1 \circ f_\nu(z)| \leq \frac{1}{\varepsilon} \rho_{\mathbb{D}^*}^{-2}(z^*) |\varphi_1(z^*)| \quad (z^* = \bar{z}^{-1})$$

for some  $\varepsilon > 0$  and for every  $z \in \mathbb{D}$ ;

(ii) the support of the complex dilatation  $\mu_0$  of the normalized quasiconformal homeomorphism  $f_0: \mathbb{D} \rightarrow \mathbb{D}$ , which is conformally conjugate to  $\widehat{f}_0: \widehat{f}_1(\Omega) \rightarrow \widehat{f}(\Omega)$ , is contained in a compact subset of  $\mathbb{D}$ ;

(iii) for the complex dilatation  $\mu_1$  of the normalized quasiconformal homeomorphism  $f_1: \mathbb{D} \rightarrow \mathbb{D}$ , which is conformally conjugate to  $\widehat{f}_1: \Omega \rightarrow \widehat{f}_1(\Omega)$ , we have

$$\varphi - \varphi_1 = \Phi(\mu_0 * \mu_1 * \nu) - \Phi(\mu_1 * \nu).$$

Combining all those maps in the claim above, we have the following commutative diagram, where  $g_\nu$ ,  $g_1$ , and  $g$  are the conjugating conformal maps:

$$\begin{array}{ccccccc}
 \mathbb{D} & \xrightarrow{f^\nu} & \mathbb{D} & \xrightarrow{f_1} & \mathbb{D} & \xrightarrow{f_0} & \mathbb{D} \\
 & \searrow f_\nu & \downarrow g_\nu & & \downarrow g_1 & & \downarrow g \\
 & & \Omega & \xrightarrow{\widehat{f}_1} & \widehat{f}_1(\Omega) & \xrightarrow{\widehat{f}_0} & \widehat{f}(\Omega) \\
 & & & \searrow \widehat{f} = \widehat{f}_0 \circ \widehat{f}_1 & & & 
 \end{array}$$

$f = f_0 \circ f_1$  (top arc)  
 $\widehat{f} = \widehat{f}_0 \circ \widehat{f}_1$  (bottom arc)

We take  $\varphi \in \mathcal{B}_0(\mathbb{D}^*)$  such that  $\varphi + \psi \in \beta(T_b)$ . Since  $\mathcal{B}_0(\mathbb{D}^*) \subset \mathcal{B}_0(\mathbb{D}^*)$ , there is a quasiconformal homeomorphism  $\widehat{f}: \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$  conformal on  $\Omega^*$  and asymptotically conformal on  $\Omega$  such that  $\mathcal{S}(\widehat{f} \circ f_\nu|_{\mathbb{D}^*}) = \varphi + \psi$ . According to the claim above, we consider the decomposition  $\widehat{f} = \widehat{f}_0 \circ \widehat{f}_1$  together with other maps that appear in it, and apply the properties shown there.

Since  $\varphi \in \mathcal{B}_0(\mathbb{D}^*)$ , if  $\varphi - \varphi_1 \in \mathcal{B}_0(\mathbb{D}^*)$ , then  $\varphi_1 \in \mathcal{B}_0(\mathbb{D}^*)$ . By property (ii),  $\mu_0$  in particular belongs to  $\mathcal{M}_0(\mathbb{D})$ , and by property (iii),  $\varphi - \varphi_1 = \Phi(\mu_0 * \mu_1 * \nu) - \Phi(\mu_1 * \nu)$ . By the previous arguments showing the inclusion  $\subset$ , we see that  $\varphi - \varphi_1 \in \mathcal{B}_0(\mathbb{D}^*)$ . Hence,  $\varphi_1 \in \mathcal{B}_0(\mathbb{D}^*)$ .

By property (i),  $\varphi_1 \in \mathcal{B}_0(\mathbb{D}^*)$  implies that  $\widehat{\mu}_1 \circ f_\nu \in \mathcal{M}_0(\mathbb{D})$ . Since  $|\widehat{\mu}_1 \circ f_\nu| = |\mu_1 \circ f_\nu|$ , we have  $\mu_1 \circ f_\nu \in \mathcal{M}_0(\mathbb{D})$ . It follows from the bi-Lipschitz continuity of  $f_\nu$  and [27, Proposition 3.5] that  $\mu_1 \in \mathcal{M}_0(\mathbb{D})$ . By property (ii), the support of the complex dilatation  $\mu_0$  of  $f_0$  is contained in a compact subset of  $\mathbb{D}$ . Hence, we see that the complex dilatation  $\mu_f = \mu_0 * \mu_1$  of  $f = f_0 \circ f_1$  belongs to  $\mathcal{M}_0(\mathbb{D})$ . Since the complex dilatation of the quasiconformal homeomorphism  $\widehat{f} \circ f_\nu$  on  $\mathbb{D}$  is  $r_\nu^{-1}(\mu_f)$ , we have that

$$\varphi + \psi = \Phi(\mu_0 * \mu_1 * \nu) = \Phi \circ r_\nu^{-1}(\mu_f) \in \Phi \circ r_\nu^{-1}(\mathcal{M}_0(\mathbb{D})) = \beta \circ R_\tau^{-1}(T_v),$$

which proves the inclusion  $\supset$ . □

## 6. Proofs of Corollaries 2 and 3

**6.1. Proof of Corollary 2.** For the quotient maps  $p: T_b \rightarrow T_v \setminus T_b$  and  $P: \mathcal{B}(\mathbb{D}^*) \rightarrow \mathcal{B}_0(\mathbb{D}^*) \setminus \mathcal{B}(\mathbb{D}^*)$ , the following commutative diagram holds:

$$\begin{array}{ccc} T_b & \xrightarrow{\beta} & \mathcal{B}(\mathbb{D}^*) \\ \downarrow p & & \downarrow P \\ T_v \setminus T_b & \xrightarrow{\widehat{\beta}} & \mathcal{B}_0(\mathbb{D}^*) \setminus \mathcal{B}(\mathbb{D}^*) \end{array}$$

The well-definedness and the injectivity of  $\widehat{\beta}$  are direct consequences from Theorem 1. Since  $P$  is the projection onto the quotient Banach space, the image  $P(\beta(T_b)) = \widehat{\beta}(T_v \setminus T_b)$  is an open subset of  $\mathcal{B}_0(\mathbb{D}^*) \setminus \mathcal{B}(\mathbb{D}^*)$ . Moreover,  $\widehat{\beta}: T_v \setminus T_b \rightarrow \widehat{\beta}(T_v \setminus T_b)$  is open and continuous because so is  $\beta: T_b \rightarrow \beta(T_b)$ . Combined with the injectivity of  $\widehat{\beta}$ , this implies that  $\widehat{\beta}$  is a homeomorphism of  $T_v \setminus T_b$  onto its image. □

**6.2. Proof of Corollary 3.** For each  $\sigma \in T_b$ , we have that  $R_\tau(T_v * \sigma) = T_v * (\sigma * \tau^{-1})$ . This shows that the correspondence  $[\sigma] \mapsto [\sigma * \tau^{-1}]$  yields a well-defined map  $\widehat{R}_\tau: p(T_b) \rightarrow p(T_b)$  that satisfies  $p \circ R_\tau = \widehat{R}_\tau \circ p$ . By considering the inverse mapping  $R_\tau^{-1} = R_{\tau^{-1}}$ , we see that  $\widehat{R}_\tau$  is bijective. In the same way as the proof of Corollary 2,  $\widehat{R}_\tau$  is shown to be a homeomorphism. Thus, it suffices to prove that  $\widehat{R}_\tau$  is holomorphic.

We may identify  $T_b$  with the domain  $\beta(T_b)$  in  $\mathcal{B}(\mathbb{D}^*)$ . The conjugate  $\widetilde{R}_\varphi = \beta \circ R_\tau \circ \beta^{-1}$  for  $\varphi = \beta(\tau)$  is a biholomorphic automorphism of  $\beta(T_b) \subset \mathcal{B}(\mathbb{D}^*)$ . We use its projection  $\widehat{R}_\varphi$  to  $P(\beta(T_b)) = \widehat{\beta}(p(T_b))$  as a replacement of  $\widehat{R}_\tau$ , which satisfies  $P \circ \widetilde{R}_\varphi = \widehat{R}_\varphi \circ P$ . Let  $\phi_1, \phi_2 \in \beta(T_b)$  with  $\phi_1 - \phi_2 \in \mathcal{B}_0(\mathbb{D}^*)$  and let  $\psi_1, \psi_2 \in \mathcal{B}(\mathbb{D}^*)$  with  $\psi_1 - \psi_2 \in \mathcal{B}_0(\mathbb{D}^*)$ . The derivative of  $\widetilde{R}_\varphi$  satisfies

$$\begin{aligned} d_{\phi_1} \widetilde{R}_\varphi(\psi_1) &= \lim_{t \rightarrow 0} \frac{1}{t} (\widetilde{R}_\varphi(\phi_1 + t\psi_1) - \widetilde{R}_\varphi(\phi_1)); \\ d_{\phi_2} \widetilde{R}_\varphi(\psi_2) &= \lim_{t \rightarrow 0} \frac{1}{t} (\widetilde{R}_\varphi(\phi_2 + t\psi_2) - \widetilde{R}_\varphi(\phi_2)), \end{aligned}$$

where the limits refer to the convergence under the norm  $\|\cdot\|_{\mathcal{B}}$ . From this, we see that  $d_{\phi_1} \widetilde{R}_\varphi(\psi_1) - d_{\phi_2} \widetilde{R}_\varphi(\psi_2)$  belongs to  $\mathcal{B}_0(\mathbb{D}^*)$  because  $\mathcal{B}_0(\mathbb{D}^*)$  is closed and

$$\begin{aligned} &\{\widetilde{R}_\varphi(\phi_1 + t\psi_1) - \widetilde{R}_\varphi(\phi_1)\} - \{\widetilde{R}_\varphi(\phi_2 + t\psi_2) - \widetilde{R}_\varphi(\phi_2)\} \\ &= \{\widetilde{R}_\varphi(\phi_1 + t\psi_1) - \widetilde{R}_\varphi(\phi_2 + t\psi_2)\} - \{\widetilde{R}_\varphi(\phi_1) - \widetilde{R}_\varphi(\phi_2)\} \end{aligned}$$

belongs to  $\mathcal{B}_0(\mathbb{D}^*)$ . Thus, for every  $[\phi] \in P(\beta(T_b))$ , a linear map  $A_{[\phi]}^\varphi: \mathcal{B}_0(\mathbb{D}^*) \setminus \mathcal{B}(\mathbb{D}^*) \rightarrow \mathcal{B}_0(\mathbb{D}^*) \setminus \mathcal{B}(\mathbb{D}^*)$  is well-defined by  $A_{[\phi]}^\varphi([\psi]) = [d_\phi \widetilde{R}_\varphi(\psi)]$ . This satisfies  $A_{[\phi]}^\varphi \circ P = P \circ d_\phi \widetilde{R}_\varphi$ .

The linear operator  $A_{[\phi]}^\varphi$  is bounded and the operator norm satisfies  $\|A_{[\phi]}^\varphi\| \leq \|d_\phi \widetilde{R}_\varphi\|$ . Indeed, for every  $[\psi] \in \mathcal{B}_0(\mathbb{D}^*) \setminus \mathcal{B}(\mathbb{D}^*)$  and every  $\varepsilon > 0$ , we may choose

$\psi \in \mathcal{B}(\mathbb{D}^*)$  such that  $P(\psi) = [\psi]$  and  $\|\psi\| \leq \|[\psi]\| + \varepsilon$ . Then,

$$\|A_{[\phi]}^\varphi([\psi])\| = \|P \circ d_\phi \tilde{R}_\varphi(\psi)\| \leq \|d_\phi \tilde{R}_\varphi(\psi)\| \leq \|d_\phi \tilde{R}_\varphi\| \cdot \|\psi\| \leq \|d_\phi \tilde{R}_\varphi\|(\|[\psi]\| + \varepsilon).$$

Making  $\varepsilon > 0$  arbitrarily small, we obtain the claim.

Moreover, since we may assume that  $\|\psi\| \leq 2\|[\psi]\|$  in the above choice of  $\psi$ , we have that

$$\begin{aligned} & \|\widehat{R}_\varphi([\phi] + [\psi]) - \widehat{R}_\varphi([\phi]) - A_{[\phi]}^\varphi([\psi])\| \\ &= \|P \circ \tilde{R}_\varphi(\phi + \psi) - P \circ \tilde{R}_\varphi(\phi) - P \circ d_\phi \tilde{R}_\varphi(\psi)\| \\ &\leq \|\tilde{R}_\varphi(\phi + \psi) - \tilde{R}_\varphi(\phi) - d_\phi \tilde{R}_\varphi(\psi)\| = o(\|[\psi]\|). \end{aligned}$$

This implies that  $\widehat{R}_\varphi$  is differentiable at every  $[\phi] \in P(\beta(T_b))$  in every direction  $[\psi] \in \mathcal{B}_0(\mathbb{D}^*) \setminus \mathcal{B}(\mathbb{D}^*)$  with the derivative  $d_{[\phi]} \widehat{R}_\varphi([\psi]) = A_{[\phi]}^\varphi([\psi])$ .  $\square$

## 7. Proof of Theorem 4

**7.1. Proof of statement (a).** Let  $V$  denote a subset of  $T_b$  consisting of all  $\tau$  for which there exists an open neighborhood  $W$  such that each  $\tau' \in W$  can be represented as a finite composition of elements in  $T_c$ . Since  $T_b$  is connected, in order to prove that  $V$  coincides with  $T_b$ , it suffices to show that  $V$  is non-empty, open, and closed. By Proposition E,  $T_c$  is an open subset of  $T_b$  containing the origin  $o = [0]$ . We see that  $o \in V$ , and hence  $V$  is non-empty. By the definition of  $V$ , this is open.

Now we prove that  $V$  is closed. Let  $\{\tau_n\} \subset V$  be a sequence such that  $\tau_n \rightarrow \tau$  as  $n \rightarrow \infty$ . We will show that  $\tau \in V$ . Let  $U$  be an open neighborhood of  $o$  in  $T_c$ . Then,  $R_\tau(\tau_n) \in U$  for all sufficiently large  $n$ , that is,  $\sigma_n = \tau_n * \tau^{-1} \in U$ . Since  $\sigma_n \rightarrow o$  as  $n \rightarrow \infty$ , we have  $\sigma_n^{-1} \rightarrow o$ . Thus, we may assume that  $\sigma_n^{-1} \in U$ . Let  $W = R_\tau^{-1}(U)$ , which is a neighborhood of  $\tau$ . For each  $\tau' \in W$ , there exists an element  $\sigma' \in U$  such that  $\tau' * \tau^{-1} = \sigma'$ . It follows that  $\tau' = \sigma' * \tau = \sigma' * \sigma_n^{-1} * \tau_n$ . Therefore,  $\tau' \in W$  can be represented as a finite composition of elements in  $T_c$ . This shows that  $\tau \in V$ .  $\square$

**7.2. Proof of statement (b).** If  $h = g^{-1} \circ f$  is the conformal welding homeomorphism corresponding to a chord-arc curve  $\Gamma$ , then  $h^{-1} = f^{-1} \circ g = (j \circ f \circ j)^{-1} \circ (j \circ g \circ j)$  is the conformal welding homeomorphism corresponding to  $j(\Gamma)$ , where  $j(z) = z^* = \bar{z}^{-1}$  is the standard reflection with respect to  $\mathbb{S}$ . Since  $j$  is an isometry with respect to the spherical metric of  $\widehat{\mathbb{C}}$ ,  $j(\Gamma)$  is a chord-arc curve. This proves that if  $\tau = [h] \in T_c$  then  $\tau^{-1} \in T_c$ .  $\square$


**7.3. Proof of statement (c).** For any  $\tau \in T_c$  and  $\sigma \in T_v$ , we will show that  $R_\tau^{-1}(\sigma)$  belongs to  $T_c$ . Set  $\widehat{\sigma} = R_\tau^{-1}(\sigma) = \sigma * \tau$ . Let  $g^{-1} \circ f$  and  $g_1^{-1} \circ f_1$  be the conformal welding homeomorphisms such that  $[g^{-1} \circ f] = \tau$  and  $[g_1^{-1} \circ f_1] = \widehat{\sigma}$ , respectively. Here,  $g$  and  $g_1$  are conformal maps on  $\mathbb{D}$ , and  $f$  and  $f_1$  are conformal maps on  $\mathbb{D}^*$  with the normalization (1). We set  $\Omega = g(\mathbb{D})$ ,  $\Omega^* = f(\mathbb{D}^*)$ , and  $\Gamma = \partial\Omega = \partial\Omega^*$  which is a chord-arc curve. Similarly, we set  $\Omega_1 = g_1(\mathbb{D})$ ,  $\Omega_1^* = f_1(\mathbb{D}^*)$ , and  $\Gamma_1 = \partial\Omega_1 = \partial\Omega_1^*$  which is a quasicircle at this moment. Let  $f^\nu$  and  $f^\mu$  be the normalized quasiconformal self-homeomorphisms of  $\mathbb{D}$  corresponding to  $\tau$  and  $\sigma$ , respectively. As  $\sigma \in T_v$ , we can assume that the complex dilatation  $\mu$  of  $f^\mu$  induces a vanishing Carleson measure  $\lambda_\mu \in \text{CM}_0(\mathbb{D})$ . Then,  $g \circ f^\nu$  and  $g_1 \circ f^\mu \circ f^\nu$  are quasiconformal extensions of  $f$  and  $f_1$  to  $\mathbb{D}$ , respectively.

We define

$$\widehat{f} = \begin{cases} f_1 \circ f^{-1} & \text{on } \Omega^*, \\ (g_1 \circ f^\mu \circ f^\nu) \circ (g \circ f^\nu)^{-1} = g_1 \circ f^\mu \circ g^{-1} & \text{on } \Omega. \end{cases}$$

Then,  $\widehat{f}: \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$  is conformal on  $\Omega^*$  and asymptotically conformal on  $\Omega$  whose complex dilatation  $\widehat{\mu}$  satisfies  $|\widehat{\mu}|^2 \rho_\Omega = \lambda_\mu \circ g^{-1} |(g^{-1})'|$  for the Poincaré density  $\rho_\Omega$  on  $\Omega$ . As  $\lambda_\mu \in \text{CM}_0(\mathbb{D})$ , we have that  $|\widehat{\mu}|^2 \rho_\Omega \in \text{CM}_0(\Omega)$  by [29, Theorem 3.2].

We decompose  $\widehat{f}$  into  $\widehat{f}_0 \circ \widehat{f}_1$  as follows. The quasiconformal homeomorphism  $\widehat{f}_1: \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$  is chosen so that its complex dilatation  $\widehat{\mu}_1$  coincides with  $\widehat{\mu}$  on  $\Omega - \Omega_0$  for some compact subset  $\Omega_0$  of  $\Omega$  homeomorphic to a closed disk, and zero elsewhere. We may assume that  $\widehat{f}_1$  satisfies the normalization (1). Then,  $\widehat{f}_0$  is defined to be  $\widehat{f} \circ \widehat{f}_1^{-1}$ . We have the following commutative diagram:



Here, the compact subset  $\Omega_0 \subset \Omega$  is chosen so that  $|\widehat{\mu}_1|^2 \rho_\Omega \in \text{CM}_0(\Omega)$  has a sufficiently small norm as a Carleson measure. It follows from [29, Lemma 4.1] that  $|\mathcal{S}(\widehat{f}_1)|^2 \rho_{\Omega^*}^{-3} \in \text{CM}_0(\Omega^*)$  with a small norm. By [29, Theorem 3.1], we have that

$$|\mathcal{S}(\widehat{f}_1 \circ f) - \mathcal{S}(f)|^2 \rho_{\mathbb{D}^*}^{-3} = (|\mathcal{S}(\widehat{f}_1)|^2 \rho_{\Omega^*}^{-3}) \circ f |f'| \in \text{CM}_0(\mathbb{D}^*),$$

and moreover, we see that it can be of a small norm according to that of  $|\mathcal{S}(\widehat{f}_1)|^2 \rho_{\Omega^*}^{-3}$ . Combined with the facts that  $\Gamma$  is a chord-arc curve and that the subspace  $T_c$  is open in  $T_b$  by Proposition E, this implies that  $\partial \widehat{f}_1(\Omega)$  is also a chord-arc curve.

Since the complex dilatation  $\widehat{\mu}_0$  of  $\widehat{f}_0$  has the compact support  $\widehat{f}_1(\Omega_0) \subset \widehat{f}_1(\Omega)$ , we conclude that  $\Gamma_1$  is the image of  $\partial \widehat{f}_1(\Omega)$  under  $\widehat{f}_0$ , which is conformal when restricted to  $\mathbb{C} - \widehat{f}_1(\Omega_0)$ . In particular,  $\widehat{f}_0$  is bi-Lipschitz on the chord-arc curve  $\partial \widehat{f}_1(\Omega)$ . Then, this extends to a bi-Lipschitz homeomorphism of  $\mathbb{C}$  (see [22, Theorem 7.10]), and thus  $\Gamma_1$  is again a chord-arc curve by Proposition D. This implies that  $\widehat{\sigma} \in T_c$ .  $\square$

*Acknowledgement.* The authors would like to thank the referee for reading the manuscript carefully and giving suggestions which improved several mathematical expressions.

## References

- [1] AHLFORS, L.: Lectures on quasiconformal mappings. - Van Nostrand, Princeton, 1966.
- [2] ASTALA, K., and M. J. GONZÁLEZ: Chord-arc curves and the Beurling transform. - Invent. Math. 205, 2016, 57–81.

- [3] ASTALA, K., and M. ZINSMEISTER: Teichmüller spaces and BMO. - *Math. Ann.* 289, 1991, 613–625.
- [4] BEURLING, A., and L. V. AHLFORS: The boundary correspondence under quasiconformal mappings. - *Acta Math.* 96, 1956, 125–142.
- [5] BISHOP, C.: A counterexample in conformal welding concerning Hausdorff dimension. - *Michigan Math. J.* 138, 1989, 233–236.
- [6] BISHOP, C., and P. JONES: Harmonic measure,  $L^2$  estimates and the Schwarzian derivative. - *J. Anal. Math.* 62, 1994, 77–113.
- [7] COIFMAN, R. R., and C. FEFFERMAN: Weighted norm inequalities for maximal functions and singular integrals. - *Studia Math.* 51, 1974, 241–250.
- [8] CUI, G., and M. ZINSMEISTER: BMO-Teichmüller spaces. - *Illinois J. Math.* 48, 2004, 1223–1233.
- [9] DOUADY, A., and C. EARLE: Conformally natural extension of homeomorphisms of the circle. - *Acta Math.* 157, 1986, 23–48.
- [10] EARLE, C. J., F. P. GARDINER, and N. LAKIC: Asymptotic Teichmüller space. Part I: The complex structure. - In: *The tradition of Ahlfors and Bers, Contemp. Math.* 256, Amer. Math. Soc., 2000, 17–38.
- [11] EARLE, C. J., V. MARKOVIC, and D. SARIC: Barycentric extension and the Bers embedding for asymptotic Teichmüller space. - In: *Complex manifolds and hyperbolic geometry, Contemp. Math.* 311, Amer. Math. Soc., 2002, 87–105.
- [12] GARDINER, F. P., and N. LAKIC: Quasiconformal Teichmüller theory. - *Math. Surveys Monogr.* 76, Amer. Math. Soc., 2000.
- [13] GARDINER, F. P., and D. SULLIVAN: Symmetric structure on a closed curve. - *Amer. J. Math.* 114, 1992, 683–736.
- [14] GARNETT, J. B.: *Bounded analytic functions.* - Academic Press, New York, 1981.
- [15] JERISON, D. S., and C. E. KENIG: Hardy spaces,  $A_\infty$ , and singular integrals on chord-arc domains. - *Math. Scand.* 50, 1982, 221–247.
- [16] LEHTO, O.: *Univalent functions and Teichmüller spaces.* - Springer-Verlag, New York, 1986.
- [17] MACMANUS, P.: Quasiconformal mappings and Ahlfors–David curves. - *Trans. Amer. Math. Soc.* 343, 1994, 853–881.
- [18] MATSUZAKI, K.: Injectivity of the quotient Bers embedding of Teichmüller spaces. - *Ann. Acad. Sci. Fenn. Math.* 44, 2019, 657–679.
- [19] MIYACHI, H.: On invariant distances on asymptotic Teichmüller spaces. - *Proc. Amer. Math. Soc.* 134, 2005, 1917–1925.
- [20] NAG, S.: *The complex analytic theory of Teichmüller spaces.* - Wiley-Interscience, New York, 1988.
- [21] POMMERENKE, CH.: On univalent functions, Bloch functions and VMOA. - *Math. Ann.* 236, 1978, 199–208.
- [22] POMMERENKE, CH.: *Boundary behaviour of conformal maps.* - Springer, 1992.
- [23] SARASON, D.: Functions of vanishing mean oscillation. - *Trans. Amer. Math. Soc.* 207, 1975, 391–405.
- [24] SEMMES, S.: A counterexample in conformal welding concerning chord-arc curves. - *Ark. Mat.* 24, 1988, 141–158.
- [25] SEMMES, S.: Quasiconformal mappings and chord-arc curves. - *Trans. Amer. Math. Soc.* 306, 1988, 233–263.
- [26] SHEN, Y., and H. WEI: Universal Teichmüller space and BMO. - *Adv. Math.* 234, 2013, 129–148.

- [27] TANG, S., H. WEI, and Y. SHEN: On Douady–Earle extension and the contractibility of the VMO-Teichmüller space. - *J. Math. Anal. Appl.* 442, 2016, 376–384.
- [28] WEI, H.: A note on the BMO-Teichmüller space. - *J. Math. Anal. Appl.* 435, 2016, 746–753.
- [29] WEI, H., and M. ZINSMEISTER: Carleson measures and chord-arc curves. - *Ann. Acad. Sci. Fenn. Math.* 43, 2018, 669–683.
- [30] WU, Y., and Y. QI: Douady–Earle extension of the strongly symmetric homeomorphism. - *Kodai Math. J.* 39, 2016, 410–424.
- [31] ZINSMEISTER, M.: *Domaines de Lavrentiev*. - *Publ. Math., Orsay*, 1985.

Received 30 April 2022 • Revision received 1 October 2022 • Accepted 3 October 2022

Published online 17 October 2022

Katsuhiko Matsuzaki  
Waseda University, School of Education  
Department of Mathematics  
Shinjuku, Tokyo 169-8050, Japan  
matsuzak@waseda.jp

Huaying Wei  
Jiangsu Normal University  
Department of Mathematics and Statistics  
Xuzhou 221116, P.R. China  
and Waseda University, School of Education  
Department of Mathematics  
Shinjuku, Tokyo 169-8050, Japan  
hywei@jsnu.edu.cn