

ON A THEOREM OF NEHARI AND QUASIDISCS

Martin Chuaqui

University of Pennsylvania, Department of Mathematics
David Rittenhouse Laboratory, 209 South 33rd Street, Philadelphia, PA 19104-6395, U.S.A.

Abstract. Let f be a locally injective analytic map of the unit disc D and let $\{f, z\}$ be its Schwarzian derivative. Suppose $|\{f, z\}| \leq 2p(|z|)$. We use the classical connection between Schwarzian derivative and second order linear equations to show that, for a particular class of functions p , the image $f(D)$ is a quasidisc. The analysis centers on the differential equation $y'' + py = 0$ and a finiteness condition of a positive solution y . The proofs are based on Sturm comparison theorems. When p in the class is analytic and $x = 1$ is a regular singular point of the linear equation, it is possible to obtain precise information about Hölder continuity of f from considerations on the Frobenius solutions at that point. The main result in this paper resolves the complementary case in a general theorem of univalence of Nehari.

1. Introduction

Let f be analytic and locally univalent, and let $\{f, z\} = (f''/f')' - \frac{1}{2}(f''/f')^2$ be its Schwarzian derivative. Two main features of the Schwarzian derivative are that all solutions to $\{f, z\} = 0$ are given by fractional linear transformations $T(z) = (az + b)/(cz + d)$ and that $\{T \circ f, z\} = \{f, z\}$. The second property is a consequence of the first and an important addition formula for the Schwarzian derivative of a composition. There is a classical connection between the Schwarzian derivative and second order linear equations: any solution of $\{f, z\} = 2p(z)$ is given by $(au + bv)/(cu + dv)$, $ad - bc \neq 0$, where u, v are two linearly independent solutions of the equation

$$(1.1) \quad y'' + py = 0.$$

A well known fact that follows from this is that f is univalent on a given domain if and only if any nontrivial solution of (1.1) vanishes at most once in the domain (see, e.g., [D]). This characterization of univalence was systematically used by Nehari, who derived several sufficient conditions for global injectivity. Briefly, estimates on the size of $|\{f, z\}|$ together with comparison theorems for the solutions of differential equations imply the absence of multiple zeroes of nontrivial solutions of (1.1). In the unit disc D , some of the conditions of this type that imply univalence are

$$(1.2) \quad |\{f, z\}| \leq \frac{\pi^2}{2},$$

$$(1.3) \quad |\{f, z\}| \leq \frac{2}{(1 - |z|^2)^2},$$

$$(1.4) \quad |\{f, z\}| \leq \frac{4}{1 - |z|^2}.$$

The constants $\pi^2/2$, 2 and 4 are sharp in each case. All of these are particular instances of the following general result due to Nehari (Theorem 1 in [N 2]):

Let $p(x) \geq 0$ be an even function on $(-1, 1)$ such that $(1 - x^2)^2 p(x)$ is nonincreasing for $x > 0$.

Suppose that the solution y of

$$(1.5) \quad y'' + py = 0, \quad y(0) = 1, \quad y'(0) = 0$$

does not vanish on $(-1, 1)$. If $|\{f, z\}| \leq 2p(|z|)$ then f is univalent in the unit disc.

With the choices for p as in (1.2), (1.3) and (1.4) the respective solutions of (1.5) are given by $\cos(\frac{1}{2}\pi x)$, $\sqrt{1 - x^2}$ and $1 - x^2$. The odd solution of the linear equation is

$$y(x) = \int_0^x y^{-2}(s) ds$$

and since the functions involved here are analytic on $(-1, 1)$ they extend to the unit disc. Consequently

$$F(z) = \int_0^z y^{-2}(\zeta) d\zeta$$

gives in each case the extremal map with the normalizations $F(0) = 0$, $F'(0) = 1$ and $F''(0) = 0$.

By using standard comparison theorems for solutions of differential equations one can go a step further and derive upper and lower bounds for $|f|$ and $|f'|$ when f is normalized as F and $|\{f, z\}| \leq 2p(|z|)$ ([C-O]).

In Nehari's theorem p is assumed to be continuous, and for such p we will continue to denote by F the associated function defined on $(-1, 1)$. Nehari also showed under what circumstances the condition $|\{f, z\}| \leq 2p(|z|)$ is sharp. It states that if $F(x) \rightarrow \infty$ as $x \rightarrow 1$ then for any positive function $r(x)$ on $(-1, 1)$ the condition

$$|\{f, z\}| \leq 2p(|z|) + r(|z|)$$

is in general not sufficient for univalence (Theorem 2 in [N 2]).

In this paper we shall be concerned with the question of what happens when $F(1) < \infty$. Our main result is

Theorem 1. *Let $p(x) \geq 0$ be an even function on $(-1, 1)$ with $(1 - x^2)^2 p(x)$ nonincreasing for $x > 0$. Suppose that the even solution y of (1.1) is positive and is such that*

$$\int_0^1 y^{-2}(x) dx < \infty.$$

If $|\{f, z\}| \leq 2p(|z|)$ then $f(D)$ is a quasidisk.

A quasidisk is the image of D under some map which is quasiconformal in the entire plane. Let Ω be simply-connected with its Poincaré metric $\lambda(z) |dz|$. A combined result of Ahlfors and later Gehring gives a characterization of quasidisks which is necessary and sufficient: there exists a positive constant η such that the inequality

$$|\{\phi, z\}| \leq \eta \lambda^2(z)$$

implies that ϕ is univalent in Ω (see, e.g., [L]).

The function p in Theorem 1 is assumed to be continuous. As previously shown, there are important cases when p is actually analytic and $x = 1$ is a regular singular point of (1.1). This allows to simplify the analysis by considering the possible Frobenius solutions at $x = 1$. The assumption that $F(1)$ is finite implies that either $y \sim (1 - x)^m$ as $x \rightarrow 1$, for some $0 < m < \frac{1}{2}$, or else that $y(1) > 0$. In the latter case, the normalized function f will be Lipschitz continuous on D while in the former case, it is possible to prove Hölder continuity.

I would like to thank C. Epstein for helpful discussions concerning the proof of Lemma 1. The referee's valuable comments allowed a simplification of the original proof and gave greater clarity to other parts of the exposition.

2. Proofs

The proof of Theorem 1 will be divided in a series of lemmas. In what follows, let p and y satisfy the hypothesis of the theorem. Let $\alpha \in [0, 1)$. Most of the analysis ahead depends on the solution u of

$$(2.1) \quad u'' + \frac{\alpha}{(1 - x^2)^2} u = 0, \quad u(0) = 1, \quad u'(0) = 0.$$

This function is given explicitly by

$$u(x) = \frac{1}{2} \sqrt{1 - x^2} \left\{ \left(\frac{1 + x}{1 - x} \right)^\beta + \left(\frac{1 - x}{1 + x} \right)^\beta \right\}$$

where $\beta = \frac{1}{2} \sqrt{1 - \alpha}$ [K, p. 492]. In particular,

$$u(x) \sim (1 - x)^{\frac{1}{2} - \beta}, \quad x \rightarrow 1,$$

and therefore $\int_0^1 u^{-2}(x) dx < \infty$. Let $\mu = \lim_{x \rightarrow 1} (1-x^2)^2 p(x)$. Clearly $\mu \geq 0$ and we claim that $\mu < 1$. If not then $p(x) \geq (1-x^2)^{-2}$. Let $P(x) = (1-x^2)^{-2}$ so that the function $q(x) = p(x) - P(x)$ is non-negative. Then $z(x) = \sqrt{1-x^2}$ satisfies

$$(2.2) \quad z'' + Pz = 0, \quad z(0) = 1, \quad z'(0) = 0.$$

Multiplying (2.2) by y , (1.5) by z , and subtracting, we get

$$z''y - zy'' = qyz.$$

We integrate this equation, using the initial condition on y and z , to obtain

$$\left(\frac{z}{y}\right)'(x) = \frac{\int_0^x (uqy)(s) ds}{y(x)^2}.$$

Hence $(z/y)'$ has the same sign as x and therefore $y \leq z$ on $(-1, 1)$ since $z(0) = y(0) = 1$. It follows that either y vanishes on $(-1, 1)$ or else $F(1) = \infty$. This contradiction proves our claim. Choose α such that $\mu < \alpha < 1$ and let now

$$q(x) = p(x) - \frac{\alpha}{(1-x^2)^2}.$$

Lemma 1. *Let $l = \liminf_{x \rightarrow 1} (1-x)(y'/y)$. Then $-\frac{1}{2} < l \leq 0$.*

Proof. Following an argument almost identical to the one given above, we can write

$$(2.3) \quad \frac{y'}{y} = \frac{u'}{u} - \frac{\int_0^x (uqy)(s) ds}{u(x)y(x)}.$$

Since $y'' = -py \leq 0$ we have $y' \leq 0$ on $(0, 1)$ because of the initial condition. Hence $l \leq 0$.

By considering the graph of the function F it follows from elementary geometry that

$$\lim_{x \rightarrow 1} \frac{1-x}{y^2} = 0.$$

Hence $(1-x)(uy)^{-1} \rightarrow 0$ as $x \rightarrow 1$.

On the other hand, the limit of $(1-x)(u'/u)$ as $x \rightarrow 1$ can be computed directly and it equals $-(\frac{1}{2} - \beta)$. This, together with equation (2.3) and the fact that $q(x) < 0$ for x sufficiently close to 1 imply the lemma.

Lemma 2. *There exists a constant M such that*

$$(2.4) \quad F(1) - F(x) \leq M \left(\frac{1-x}{y^2} \right).$$

Proof. The derivative of the left hand side of (2.4) is $-y^{-2}$ while the derivative of $(1-x)y^{-2}$ is

$$-y^{-2} \left(1 + 2(1-x) \frac{y'}{y} \right).$$

Lemma 1 implies that $1 + 2(1-x)(y'/y) \geq \sigma > 0$ provided x is sufficiently close to 1. Hence for all such x

$$F(1) - F(x) \leq \frac{1}{\sigma} \left(\frac{1-x}{y^2} \right)$$

and the lemma follows.

Now we state the key result in this chain.

Lemma 3. *There exists a constant $\eta > 0$ such that the solution φ of*

$$(2.5) \quad \varphi'' + \left(p(x) + \frac{\eta}{(1-x^2)^2} \right) \varphi = 0, \quad \varphi(0) = 1, \quad \varphi'(0) = 0$$

does not vanish on $(-1, 1)$.

Proof. Let $c = F(1)$. On the image interval $(-c, c)$ we consider the ‘‘Poincaré density’’

$$\lambda(w) = \frac{1}{F'(x)(1-x^2)} = \frac{y^2}{1-x^2}$$

where $w = F(x)$. We will show that for $\eta > 0$ sufficiently small the solution h of

$$(2.6) \quad h'' + \eta \lambda^2(w) h = 0, \quad h(0) = 1, \quad h'(0) = 0$$

is positive on $(-c, c)$. By Lemma 2,

$$\lambda(w) = \frac{y^2}{1-x^2} \leq \frac{y^2}{1-x} \leq \frac{M}{c-w} \leq \frac{2Mc}{c^2-w^2}.$$

Thus it suffices to show that the solution of (2.6) with $4M^2c^2(c^2-w^2)^{-2}$ instead of $\lambda^2(w)$ does not vanish. This will be the case as long as $4M^2\eta \leq 1$. To see this, we rescale. The function $\bar{h}(x) = h(cx)$ solves $\bar{h}'' + 4M^2\eta(1-x^2)^{-2}\bar{h} = 0$ with even initial conditions. Then $\bar{h} > 0$ on $(-1, 1)$ if and only if $4M^2\eta \leq 1$ [K, p. 492]. With h the positive solution of (2.6) we define φ by

$$\varphi(x) = y(x)h(F(x)).$$

A straightforward computation shows that φ is the solution of (2.5). This finishes the proof of Lemma 3.

This lemma together with Nehari's first theorem shows that

$$(2.7) \quad |\{g, z\}| \leq 2 \left(p(|z|) + \frac{\eta}{(1 - |z|^2)^2} \right)$$

is a sufficient condition for univalence. The proof of Theorem 1 is now quite simple. Assume $|\{f, z\}| \leq 2p(|z|)$ and let $\lambda(\zeta) |d\zeta|$ be the Poincaré metric on $\Omega = f(D)$. We will show that

$$|\{\phi, \zeta\}| \leq 2\eta\lambda^2(\zeta)$$

implies the univalence of the map ϕ . Let $g(z) = \phi(f(z))$. Then

$$\{g, z\} = \{\phi, f(z)\}f'(z)^2 + \{f, z\}$$

and therefore

$$(1 - |z|^2)^2 |\{g, z\}| \leq \lambda^{-2}(\zeta) |\{\phi, \zeta\}| + 2(1 - |z|^2)^2 p(|z|)$$

where $\zeta = f(z)$. It follows that g satisfies (2.7), hence g and consequently ϕ are univalent. This shows that Ω is a quasidisc.

3. The analytic case

In this section we shall assume that, in addition, p is analytic and that $x = 1$ is a regular singular point of the equation (1.1). The assumptions on the even solution y are as before. Recall that $\mu = \lim_{x \rightarrow 1} (1 - x^2)^2 p(x)$. From the analysis of the possible Frobenius solutions at $x = 1$ we will prove Hölder or Lipschitz continuity for maps f that satisfy $|\{f, z\}| \leq 2p(|z|)$. Because of the invariance of the Schwarzian derivative under Möbius changes one can not expect such a result unless f is properly normalized. The right normalization turns out to be $f''(0) = 0$. Let u solve

$$u'' + \frac{1}{2}\{f, z\}u = 0, \quad u(0) = 1, \quad u'(0) = 0$$

and let

$$v(z) = u(z) \int_0^z u^{-2}(\zeta) d\zeta$$

be the solution with odd initial conditions. If $f''(0) = 0$ then

$$f(z) = f(0) + f'(0) \int_0^z u^{-2}(\zeta) d\zeta.$$

From Lemma 2 in [C-O] it follows that if $|\{f, z\}| \leq 2p(|z|)$ then

$$|u(z)| \geq y(|z|)$$

and therefore

$$|f'(z)| \leq |f'(0)|y^{-2}(|z|).$$

We distinguish the cases $\mu > 0$ and $\mu = 0$. Suppose μ is positive. Then $p(x) \geq \mu(1-x^2)^{-2}$ and hence the function y must vanish at $x = 1$. The possible orders m of vanishing are given by the roots of the indicial equation

$$m^2 - m + \frac{\mu}{4} = 0,$$

i.e.,

$$(3.1) \quad m_1 = \frac{1 + \sqrt{1 - \mu}}{2}, \quad m_2 = \frac{1 - \sqrt{1 - \mu}}{2}.$$

Note that $0 < m_2 < \frac{1}{2} < m_1 < 1$. Since $m_1 - m_2$ is not an integer both orders of vanishing can occur [H].

Theorem 2. *Let f satisfy $|\{f, z\}| \leq 2p(|z|)$, $f''(0) = 0$ and suppose $\mu > 0$. If $F(1)$ is finite then f is Hölder continuous on D with Hölder exponent $\sqrt{1 - \mu}$.*

Proof. The assumption that $F(1) < \infty$ implies that $y \sim (1-x)^{m_2}$ as $x \rightarrow 1$. Therefore

$$|f'(z)| = O((1 - |z|)^{-2m_2}).$$

A standard technique of integrating along hyperbolic segments (see, e.g., [G-P]) gives

$$|f(z_1) - f(z_2)| = O(|z_1 - z_2|^{1-2m_2}),$$

and the theorem follows.

Suppose now $\mu = 0$. In this case, the roots of the indicial equation are 1 and 0. Hence two linearly independent solutions are $y_1 = (1-x)h_1$ and $y_2 = h_2 + cy_1 \log(1-x)$, where h_1, h_2 are analytic and nonvanishing at $x = 1$ [H, Theorem 5.3.1].

Theorem 3. *Let f satisfy $|\{f, z\}| \leq 2p(|z|)$, $f''(0) = 0$ and suppose $\mu = 0$. If $F(1)$ is finite then f is Lipschitz continuous on D .*

Proof. The finiteness condition and the discussion preceding the theorem imply that in fact y cannot vanish at $x = 1$. Hence $|f'|$ is uniformly bounded.

The following situation describes accurately the case $\mu = 0$. For $0 \leq t < 1$ let $p_t(x) = tp(x)$, where $p(x) = 2(1-x^2)^{-1}$. Since the inequality $|\{f, z\}| \leq 2p(|z|)$ is sufficient for univalence then $|\{f, z\}| \leq 2p_t(|z|)$ implies that $f(D)$ is a quasidisk (Theorem 6 in [G-P]). As mentioned in the introduction, the even solution of (1.1)

is in this case $y = 1 - x^2$. We claim that the even solution y_t of (1.1) with p replaced by p_t must be positive at the endpoints. To show this, let

$$F_t(z) = \int_0^z y_t^{-2}(\zeta) d\zeta.$$

This function is odd and has Schwarzian derivative equal to $2p_t(z)$. Therefore $F_t(D)$ is a quasidisc and hence $F_t(1) < \infty$, otherwise the point at infinity would be a point of self-intersection of $\partial F_t(D)$. This in turn would contradict the fact that $\partial F_t(D)$ is a Jordan curve. Since $\mu_t = t\mu = 0$ it follows that y_t is a linear combination of the functions y_1, y_2 as in the paragraph preceding the statement of Theorem 3. Thus $F_t(1) < \infty$ forces $y_t(1) > 0$.

We consider finally examples for any $\mu \in (0, 1)$. For $s \in (1, 2)$ let

$$p(x) = s \frac{1 - (s-1)x^2}{(1-x^2)^2}.$$

Then $\mu = s(2-s)$ and the even solution of (1.1) is

$$y = (1-x^2)^{s/2}.$$

(The exponent $s/2$ corresponds to m_1 in (3.1) and $m_2 = (2-s)/2$.) This shows that the function F has $F(1) = \infty$. On the other hand, by the argument given above, changing p to tp has the effect of making $F_t(1)$ finite. Consider now equation (3.1) with μ replaced by $t\mu$. Since $y_t \sim (1-x)^{m_1}$, $x \rightarrow 1$, would make $F_t(1)$ infinite, we conclude that the order of vanishing of the solution y_t must be the other root, m_2 .

References

- [A-H] ANDERSON, J.M., and A. HINKKANEN: Positive superharmonic functions and the Hölder continuity of conformal mappings. - J. London Math. Soc. 39, 1989, 256–270.
- [C-O] CHUAQUI, M., and B. OSGOOD: Sharp distortion theorems associated with the Schwarzian derivative. - Proc. London Math. Soc. (to appear).
- [D] DUREN, P.: Univalent functions. - Springer-Verlag, New York, 1983.
- [G-P] GEHRING, F.W., and CH. POMMERENKE: On the Nehari univalence criterion and quasi-circles. - Comment. Math. Helv. 59, 1984, 226–242.
- [H] HILLE, E.: Ordinary differential equations in the complex domain. - John Wiley & Sons, New York, 1976.
- [K] KAMKE, E.: Differentialgleichungen. - Chelsea, New York, 1948.
- [L] LEHTO, O.: Univalent functions and Teichmüller spaces. - Springer-Verlag, New York, 1987.
- [N 1] NEHARI, Z.: The Schwarzian derivative and schlicht functions. - Bull. Amer. Math. Soc. 55, 1949, 545–551.
- [N 2] NEHARI, Z.: Univalence criteria depending on the Schwarzian derivative. - Illinois J. Math. 23, 1979, 345–351.

Received 24 February 1992