

# ERGODICITY OF SOME CLASSES OF MEROMORPHIC FUNCTIONS

Linda Keen and Janina Kotus

CUNY Lehman College, Mathematics Department  
Bronx, NY 10468, U.S.A.; linda@alpha.lehman.cuny.edu  
Polish Academy of Sciences, Institute of Mathematics  
PL-00-950 Warsaw, Poland

**Abstract.** We consider the class of meromorphic functions whose set of singular values is bounded and for which the  $\omega$ -limit set of the post singular set is a compact repeller. We show that if two simple growth conditions are satisfied, then the function is ergodic on its Julia set.

## 1. Introduction

The dynamical phenomena of systems generated by conformal mappings of the plane are controlled by the behavior of the forward orbits of the singular values. The closure of these orbits is called the post-singular set, or in the case of rational maps, the post-critical set. For rational maps, it is known that outside the post-critical set the function is expanding and this expansion leads to the following dichotomy: a rational map either acts ergodically with respect to Lebesgue measure on the sphere and the Julia set is the full sphere or the postcritical set behaves as a measure theoretic attractor. (See [16], [18]). For transcendental meromorphic functions, the essential singularities make the situation more complicated. For example, in [15] Lyubich proved that for the exponential function, the set of points whose  $\omega$ -limit set contains the  $\omega$ -limit set of the post-singular set may have positive but not full measure and in [17] McMullen proved that for the sine family the set of points attracted to infinity always has positive measure and many ergodic components.

In this paper, motivated by our study of the tangent family [14], we find sufficient conditions for a transcendental meromorphic function to be ergodic on its Julia set. We prove

**Theorem 1.** *Let  $f$  be a transcendental meromorphic function whose singular values lie in a bounded set. If  $f$  satisfies two simple growth conditions, if the  $\omega$ -limit set of the post singular set,  $\omega_f$ , is compact and if for some  $k > 1$ ,  $|f'| > k$  on  $\omega_f$ , then  $f$  is ergodic with respect to Lebesgue measure on its Julia set.*

---

1991 Mathematics Subject Classification: Primary 30D05, 58F08.

The first author was supported in part by NSF and PSC-CUNY and the second author was supported in part by Polish KBN Grant No 2 P03A 025 12 “Iteracje funkcji holomorficznych”. The second author’s permanent address: Technical University of Warsaw, PL-00-661 Warsaw, Poland.

As examples of the theorem we see first that if the omitted values of the tangent map  $\lambda \tan(z)$  land on repelling periodic cycles, the map is ergodic and second that there are values of  $\lambda$  such that the map  $\lambda e^{-z^2} \sin(z)$  is ergodic. These special cases also follow from the results in [8].

Our techniques are an adaptation of those developed in [11] to control the set attracted to the essential singularities. The new ideas involve dealing with the poles.

The paper is organized as follows. In Section 2 we set our notation and summarize the basic definitions and theory. In Section 3 we prove our theorem and in Section 4 we discuss applications and examples.

We would like to thank I.N. Baker for helpful comments on the original version and for pointing out that our proof of the main theorem holds more generally than we had originally stated it.

## 2. Preliminaries

**2.1. Julia sets of meromorphic functions.** If  $f: \mathbf{C} \rightarrow \widehat{\mathbf{C}}$  is a transcendental meromorphic function, the orbits of points fall into three categories: they may be infinite, they may become periodic and hence consist of a finite number of distinct points, or they may terminate at a pole of  $f$ . To study the dynamics, we define the *stable set*, or *Fatou set*,  $\Omega_f$  as the set of those points  $z$  such that the sequence  $f^n(z)$  is defined and meromorphic for all  $n$  and forms a normal family in a neighborhood of  $z$ . The *unstable set*, or *Julia set*,  $J_f$  is the complement of the stable set. We assume  $\infty \in J_f$ . Thus,  $\Omega_f$  is open,  $J_f$  is closed. It is easy to see that  $\Omega_f$  is completely invariant and  $z \in J_f$  if and only if  $f(z) \in J_f$  or  $z = \infty$ . As in the rational case  $J_f \subset \overline{\bigcup_{n \geq 0} f^{-n}(z)}$  for all  $z \in \widehat{\mathbf{C}} - E_f$ , where  $E_f$  consists of at most two *exceptional values* with finite inverse orbits. For all transcendental entire functions  $\infty$  is an omitted value, so it is also exceptional. For generic transcendental meromorphic functions, the set of prepoles  $\mathcal{P}_f = \bigcup_{n \geq 0} f^{-n}(\infty)$  is infinite and  $J_f = \overline{\mathcal{P}_f}$ . Moreover, the Julia set  $J_f$  is the closure of the repelling periodic points ([1], [3], [9]).

The *singular set*  $S = S_f$  of a meromorphic function  $f$  consists of those values in  $\mathbf{C}$  at which  $f$  is not a regular covering. Therefore at a singular value  $v$  there is a branch of the inverse which is not holomorphic but has an algebraic or transcendental singularity. If the singularity is algebraic,  $v$  is a critical value, whereas if it is transcendental, there is a path  $\alpha: [0, \infty) \rightarrow \mathbf{C}$  such that  $\lim_{t \rightarrow \infty} |\alpha(t)| = \infty$  and  $\lim_{t \rightarrow \infty} f(\alpha(t)) = v$ , and  $v$  is called an *asymptotic value* for  $f$ . If we can associate to a given asymptotic value  $v$  an *asymptotic tract*, that is, a simply connected unbounded domain  $A$  such that  $f(A)$  is a punctured neighborhood of  $v$  and  $f: A \rightarrow f(A)$  is an unramified covering, then  $v$  is called a *logarithmic*

*singularity*. We define the *post-singular set* as

$$PS = PS_f = \overline{\bigcup_{n \geq 0} f^n(S)}$$

and denote its  $\omega$ -limit set by  $\omega_f$ .

We distinguish the following classes of meromorphic functions. For  $f$  meromorphic and  $p \geq 1$  define

$$\mathcal{M}_p = \{f : \text{card } S = p\}, \quad \mathcal{M} = \bigcup_{p \geq 1} \mathcal{M}_p, \quad \text{and} \quad \mathcal{B} = \{f : S \text{ is bounded}\}.$$

Of course  $\mathcal{M} \subset \mathcal{B}$ . Some basic examples of entire functions in  $\mathcal{M}$  are:

$$\lambda e^z, \quad a \sin(z) + b, \quad f(z) = \int^z h(\eta) \exp(p(\eta)) d\eta,$$

where  $\lambda, a, b \in \mathbf{C}$  and  $h(\eta), p(\eta)$  are polynomials. The class  $\mathcal{M}$  also includes meromorphic functions with polynomial Schwarzian derivative. An example is  $\lambda \tan(z)$  whose Schwarzian derivative is constant. Examples of functions in  $\mathcal{B} - \mathcal{M}$  are:

$$f(z) = \lambda \sin(z)/z \quad \text{and} \quad f(z) = \lambda e^{-z^2} \sin(z);$$

an example of a function not in  $\mathcal{B}$  is  $f(z) = \lambda z \sin(z)$ .

We call the set  $\omega_f$  a *repeller* if, for all  $z \in \omega_f$  such that  $f(z)$  is defined,  $f(z) \in \omega_f$  and if there exists  $k > 1$  such that, for all  $z \in \omega_f$ ,  $|f'(z)| \geq k$ . It is a *compact repeller* if  $f(z)$  is defined for all  $z \in \omega_f$  and if  $\omega_f$  is compact.

In particular, if  $f \in \mathcal{M}$  and all critical and asymptotic values are eventually mapped to repelling periodic points, then  $\omega_f$  is a compact repeller. Note that for transcendental meromorphic functions with poles,  $\omega_f$  may include points with finite forward orbits. For example, the singular set  $S$  of the function  $f(z) = \frac{1}{2}\pi i \tan(z)$  consists of the omitted values  $\{\pm \frac{1}{2}\pi\}$ . Since they are poles and have no forward orbit,  $\omega_f = S_f$  is a repeller but is not a compact repeller. Clearly, if  $\omega_f$  is not finite and contains prepoles it cannot be a compact repeller.

**2.2. Classification of stable behavior.** Let  $D$  be a component of the stable set;  $f$  will map  $D$  to a component, but if the image contains an asymptotic value, the map might not be onto. In any case, we call the image  $f(D)$  and note that either there exist integers  $m \neq n > 0$  such that  $f^n(D) = f^m(D)$ , and  $D$  is called *eventually periodic* with period  $p = \min(|m - n|)$ , with the minimum taken over all such  $m, n$ , or for all  $m \neq n$ ,  $f^n(D) \cap f^m(D) = \emptyset$ , and  $D$  is called a *wandering domain*.

The qualitative and quantitative description of the eventually periodic behavior is slightly more complicated than in the rational case because of the transcendental singularity at  $\infty$  and the possibility that  $f^p$  may not be defined at

some values. As for rational maps, eventually periodic domains may be attracting, parabolic or rotation domains. In addition, however, an eventually periodic domain  $D$  may be an *essentially parabolic or Baker* domain; that is, the boundary of  $D$  contains a point  $z_0$  (possibly  $\infty$ ) such that  $f^{np}(z) \rightarrow z_0$  for  $z \in D$  and  $f^p$  is not holomorphic at  $z_0$ . If  $p = 1$ , then the only possible boundary point is  $\infty$ .

**2.3. Two propositions.** We shall need the following proposition proved in [13].

**Proposition 2.1.** *Let  $f$  be a meromorphic function and let  $D$  be an open disk such that  $D \cap J_f \neq \emptyset$ . Suppose that there are branches  $g_{n_k} \in f^{-n}$  holomorphic and univalent on  $D$  with  $n_k \rightarrow \infty$ . Then  $(g'_{n_k}) \rightarrow 0$  uniformly on every compact set in  $D$ .*

We shall also need the following version of Koebe's distortion theorem.

**Proposition 2.2.** *If  $g: D(0, 1) \rightarrow \mathbf{C}$  is a univalent map normalized so that  $g(0) = 0$ , then for every  $0 < k < 1$  and  $z, w \in D(0, k)$ , there is a constant  $T(k)$  such that  $|g'(z)|/|g'(w)|$  lies between  $1/T(k)$  and  $T(k)$ .*

**2.4. Orbits that tend to infinity.** In this section we adapt the discussion of Eremenko and Lyubich [11] on entire functions in  $\mathcal{B}$  to meromorphic functions in  $\mathcal{B}$ . They give a sufficient condition for the measure of the set

$$I_\infty(f) = \{z \in \mathbf{C} : f^n(z) \rightarrow \infty\}$$

to be zero. We need additional conditions on the Laurent expansions about the poles to prove this in the meromorphic case.

At each pole  $p$  of order  $m_p$ , form the Laurent expansion of  $f$  about  $p$ ,  $f_p(z) = c_p/(z-p)^{m_p}(1 + \phi_p(z-p))$  where  $\phi(z)$  is analytic and  $\phi(z-p) = o((z-p)^{m_p})$ .

Suppose  $f \in \mathcal{B}$ . Let  $D(q, r)$  denote the disk of radius  $r$  centered at  $q$ ;  $A_r$  the annulus  $A_r = \{z : r < |z| < \infty\}$ .

Because  $S(f)$  is bounded, we can find  $R_0 > 0$  such that  $S(f) \subset D(0, R_0)$ . Fix  $R > R_0$ . Without loss of generality we may assume that  $f$  is analytic at 0 and  $|f(0)| < \frac{1}{2}R_0$ .

Suppose  $w \in A_R$  and  $f(z) = w$ . Since there are no singular values in  $A_R$ , any branch  $g$  of  $f^{-1}$  such that  $g(w) = z$  can be continued analytically throughout  $A_R$ . Let  $g$  be some branch of  $f^{-1}$  and set  $V = g(A_R)$ . By [20],  $\infty$  is either a logarithmic branch point of  $g$  with asymptotic value  $a$ ,  $V = V_a$  is simply connected and  $f: V_a \rightarrow A_R$  is a universal covering, or  $\infty$  is an algebraic branch point of  $g$  of order  $m_p - 1$  for some integer  $m_p$ ,  $V = V_p$  is conformally equivalent to a disk punctured at a pole  $p$  of  $f$  and  $f: V_p \rightarrow A_R$  is a regular  $m_p$  to 1 covering.

Let  $m$  denote Lebesgue measure in  $\mathbf{C}$  and let  $\Theta_R(r, f)$  be the linear measure of the set  $\{\theta : |f(re^{i\theta})| < R\}$ .

With this notation we have

**Proposition 2.3.** *Let  $f \in \mathcal{B}$  and suppose that there is a positive integer  $J$  and positive constants  $b, B, C_1, C_2 > 0$  such that for every  $p$ , and  $z \in V_p$ , the multiplicities  $m_p$  are bounded by  $J$  and the coefficients  $c_p$  and functions  $\phi_p(z - p)$  and  $\phi'_p(z - p)$  satisfy*

$$(*) \quad b < |c_p| < B, \quad |\phi_p(z - p)| < C_1, \quad |\phi'_p(z - p)| < C_1, \quad C_2 < |1 + \phi_p(z - p)|.$$

Moreover suppose

$$(**) \quad \liminf_{r \rightarrow \infty} \frac{1}{\log r} \int_1^r \Theta_R(t, f) \frac{dt}{t} > 0.$$

Then  $m(I_\infty(f)) = 0$ .

*Proof.* We summarize the proof in [11] that  $m(I_\infty) = 0$  for entire functions satisfying condition (\*\*), indicating the necessary changes to account for the poles. We remark that the proof there is valid if there are only finitely many poles. We therefore assume here that our functions have infinitely many poles.

The first step in the proof that  $m(I_\infty) = 0$  is to obtain a uniform expansion estimate on  $|f'|$  restricted to each component  $V_a$  or  $V_p$ .

For simply connected  $V_a$ , the argument is the same as in [11] using Teichmüller's construction of a logarithmic coordinate. We outline that argument here. If  $\infty$  is a logarithmic branch point of  $g$ , we first define a logarithmic coordinate  $w = \log z$ ,  $z \in V_a$  and set  $\mathcal{U}_a = \log V_a$ ; then  $\mathcal{U}_a \subset H = \{w : \operatorname{Re} w > 0\}$  has infinitely many simply connected unbounded components  $U_{a,n}$  each contained in a vertical strip of width  $2\pi$  and the exponential maps each  $U_{a,n}$  univalently onto  $V_a$ . We have a commutative diagram:

$$\begin{array}{ccc} \mathcal{U}_a & \xrightarrow{F} & H_R \\ \downarrow \exp & & \downarrow \exp \\ V_a & \xrightarrow{f} & A_R \end{array}$$

where  $H = \{w : \operatorname{Re} w > \log R\}$  and  $F$  maps  $U_{a,n} \subset H$  univalently onto  $H_R$ .

If  $w \in H$ , the disk  $D = D(F(w), \delta(w))$  centered at  $F(w)$  with radius  $\delta(w) = \operatorname{Re} F(w) - \log R$  is contained in  $H_R$ . Denote the branches of  $F^{-1}$  by  $G_n: H_R \rightarrow U_{a,n}$ . By the Koebe  $\frac{1}{4}$ -theorem  $G_n(D)$  contains a disk  $D_n$  of center  $w$  and radius  $\frac{1}{4}\delta(w)|(G_n)'(F(w))|$ . Since  $U_{a,n}$  cannot contain any vertical segment of length  $2\pi$ , for  $w \in U_{a,n}$  we have

$$|F'(w)| > \frac{1}{4\pi} (\operatorname{Re} F(w) - \log R).$$

It follows there are constants  $k, K > 1$  such that if  $\operatorname{Re} F(w) > k$ , then  $|F'| > K$ . By the chain rule, we have  $|f'(z)| > K|f(z)|/|z|$ .

To obtain an estimate for  $|f'| = |f'_p|$  on the punctured disk domain  $V_p$  we use a different argument.

Note that since  $\phi_p$  and  $\phi'_p$  are analytic in  $V_p$ ,  $|\phi_p|$  and  $|\phi'_p|$  achieve their maxima on  $\partial V_p$ . Increasing  $R$  clearly decreases these maxima so that by taking  $R$  large enough, we may assume  $C_1$ ,  $C_2$  and  $C_1/C_2$  are small relative to  $R$ .

If  $z \in V_p$ , by conditions (\*)

$$R \leq |f_p(z)| = \left| \frac{c_p}{(z-p)^{m_p}} (1 + \phi_p(z-p)) \right| \leq \frac{B}{|z-p|^{m_p}} (1 + C_1)$$

so that  $|z-p| < M_p = (B(1+C_1)/R)^{1/m_p} < 1$ .

For  $t > R$ , set  $A_{R,t} = A_R \setminus A_t$ . Choose  $R' \gg R$  and such that  $R' > C_2 b$ ; for each pole  $p$  set  $\tilde{V}_p = f_p^{-1}(A_{R,R'})$ . Then if  $z \in \tilde{V}_p$ , by conditions (\*) we have

$$\frac{bC_2}{|z-p|^{m_p}} \leq |f_p(z)| = \left| \frac{c_p}{(z-p)^{m_p}} (1 + \phi_p(z-p)) \right| \leq R'$$

so that  $|z-p| > \varrho_p = (bC_2/R')^{1/m_p}$ . The annulus  $\varrho_p < |z-p| < M_p$  is thus contained in  $\tilde{V}_p$ . Note that since for all  $p$ ,  $1 \leq m_p \leq J$ , the moduli of these annuli are uniformly bounded.

Next compute that

$$|f'_p(z)| = |f_p(z)| \left| \frac{m_p}{z-p} + \frac{\phi'_p(z-p)}{1 + \phi_p(z-p)} \right|.$$

In  $\tilde{V}_p$  we have

$$|f'_p(z)| \leq R' \left| J \left( \frac{R'}{bC_2} \right)^{1/m_p} + \frac{C_1}{C_2} \right|$$

and

$$\left| \frac{m_p}{z-p} + \frac{\phi'_p(z-p)}{1 + \phi_p(z-p)} \right| > \left| \left| \frac{m_p}{z-p} \right| - \left| \frac{\phi'_p(z-p)}{1 + \phi_p(z-p)} \right| \right| \geq \left| \frac{1}{M_p^{m_p}} - \frac{C_1}{C_2} \right|.$$

Thus by choosing  $R$  and  $R'$  large enough, we see that there are constants,  $K'$ ,  $K$ , independent of  $p$ , such that for every  $|p| > R$ , in  $\tilde{V}_p$  we have

$$(1) \quad K' > |f'_p(z)| > K > 1.$$

To show that  $m(I_\infty) = 0$ , by the Lebesgue density theorem it is sufficient to show that for any point  $z \in I_\infty$ ,

$$(2) \quad \limsup_{\delta \rightarrow 0} \frac{m(D(z, \delta) \cap I_\infty)}{m(D(z, \delta))} < 1.$$

In order to do this we first restate condition (\*\*) as follows: Let  $\mathcal{V} = f^{-1}(A_R)$ . Then there is a  $\kappa > 0$  such that

$$(3) \quad \limsup_{t \rightarrow \infty} \frac{\text{area}(A_{R,t} \cap \mathcal{V})}{\text{area}(A_{R,t})} \leq 1 - \kappa.$$

In terms of the logarithmic coordinates  $\log z = s + i\theta$ , let  $S_t = \{\log R < s < t, 0 \leq \theta < 2\pi\}$  and set  $\mathcal{U} = \log \mathcal{V}$ . Then there is a  $\delta > 0$  such that

$$(4) \quad \limsup_{t \rightarrow \infty} \frac{\text{area}(S_t \cap \mathcal{U})}{\text{area}(S_t)} \leq 1 - \delta.$$

For  $z_0 \in I_\infty$ , set  $z_n = f^n(z_0)$ . We may assume without loss of generality that  $|z_n| > R$  is so large that the estimates (1) on  $|f'|$  hold. Then either there is some  $N$  such that for all  $n > N$ ,  $z_n \in V_a$  for some simply connected component  $V_a$  or there is a subsequence  $z_{n_j} \rightarrow \infty$  and a sequence of poles  $p_j \rightarrow \infty$  such that  $z_{n_j} \in V_{p_j}$ .

In the first case the argument is just as in [11]. For  $w_0 = \log z_0$  in a fixed strip of width  $2\pi$ , let  $w_n = F(w_{n-1})$ , where  $F$  is the lift defined above. We may assume that for all  $n$ ,  $\text{Re } w_n$  is large enough that  $|F'(w_n)| > K > 1$ . Let  $F_n^{-1}(w_n) = w_{n-1}$ . Then, there is a constant  $d$ , independent of  $n$  such that  $F_n^{-1}$  is univalent on  $D_n = D(w_n, d)$ . Moreover, by the expansion of  $F$ , we may choose  $m$  such that  $F(D_n)$  contains a vertical segment of width bigger than  $2\pi$ . Fix such an  $m$  and let  $B_j = F^{-j}(D(w_m, d/4))$  for  $j = 1, \dots, m$ , where  $F^{-j}$  is the composition of the appropriate inverse branches. Applying the Koebe distortion theorem to the function  $F^{-m}$  we have  $B_m \subset D(w_0, d/4K^m)$ . Moreover if  $s_m$  is the radius of the smallest disk centered at  $w_0$  containing  $B_m$ , there is a constant  $t < 1$ , independent of  $m$  such that  $D(w_0, ts_m) \subset B_m \subset D(w_0, s_m)$ . Clearly  $s_m \rightarrow 0$  as  $m \rightarrow \infty$  so that applying (4) we obtain (2).

In the second case we need to modify the argument. Since the moduli of the annuli  $\tilde{V}_p$  have the same bounds, there is a constant  $0 < \tau < 1$ , independent of  $V_p$ , such that

$$\frac{\text{area}(\tilde{V}_p)}{\text{area}(V_p)} \leq 1 - \left( \frac{bC_2R}{B|1 + C_1|R'} \right)^{2/m_p} < 1 - \tau.$$

We want to show that there is a constant  $\mu > 0$  independent of  $p$  satisfying

$$(5) \quad \frac{\text{area}(V_p \cap f_p^{-1}(\mathcal{V}))}{\text{area}(V_p)} \leq 1 - \mu.$$

With the constants of inequality (1), set  $L = K'/K$ . We claim that  $\mu = \kappa\tau/L^2$  where  $\kappa$  is the constant in inequality (3).

We rewrite  $\text{area}(V_p \cap f_p^{-1}(\mathcal{V}))/\text{area}(V_p)$  and estimate

$$\begin{aligned}
& \frac{\text{area}([(V_p \setminus \tilde{V}_p) \cap f_p^{-1}(\mathcal{V})] \cup [\tilde{V}_p \cap f_p^{-1}(\mathcal{V})])}{\text{area}(V_p)} \\
& \leq \frac{\text{area}([(V_p \setminus \tilde{V}_p) \cap f_p^{-1}(\mathcal{V})])}{\text{area}(V_p)} + \frac{\text{area}(\tilde{V}_p)}{\text{area}(V_p)} \\
& = \frac{\text{area}([(V_p \setminus \tilde{V}_p) \cap f_p^{-1}(\mathcal{V})])}{\text{area}(V_p \setminus \tilde{V}_p)} \times \frac{\text{area}(V_p \setminus \tilde{V}_p)}{\text{area}(V_p)} + \frac{\text{area}(\tilde{V}_p)}{\text{area}(V_p)} \\
& \leq \left(1 - \frac{\kappa}{L^2}\right) \left(1 - \frac{\text{area}(\tilde{V}_p)}{\text{area}(V_p)}\right) + \frac{\text{area}(\tilde{V}_p)}{\text{area}(V_p)} \\
& = 1 - \frac{\kappa}{L^2} - \frac{\text{area}(\tilde{V}_p)}{\text{area}(V_p)} + \frac{\kappa}{L^2} \frac{\text{area}(\tilde{V}_p)}{\text{area}(V_p)} + \frac{\text{area}(\tilde{V}_p)}{\text{area}(V_p)} \\
& \leq 1 - \frac{\kappa}{L^2} + \frac{\kappa}{L^2}(1 - \tau) = 1 - \frac{\kappa}{L^2}\tau = 1 - \mu.
\end{aligned}$$

Now fix  $n_j$  such that  $z_{n_j} \in V_{n_j}$  and choose a univalent branch of  $f_{p_{n_j-1}}^{-1}$  so that  $f_{p_{n_j-1}}^{-1}: z_{n_j} \mapsto z_{n_j-1}$ . Let  $m = n_j$  and set  $D_m = D(z_m, M_{p_m})$ . For large  $m$ ,  $|z_m - p_m| \ll \varrho_{p_m}$  and  $V_{p_m} \approx D(z_m, M_{p_m})$ . We define  $B_m = f^{-m}(z_m, M_{p_m}/4)$  where again  $f^{-m}$  is a composition of the appropriate inverse branches so that  $z_0 \in B_m$ . Then  $f^{-m}$  is univalent on  $V_{p_m}$  and we can control distortion on  $D_m$ . Now for any trajectory from  $z_n$  to  $z_{n+k}$ , by the chain rule and the estimates on factors  $|f'|$ , whether the corresponding domain containing  $z_j$  is a simply connected  $V_a$  or is an annular  $\tilde{V}_p$ , we have  $|f^k(z_{n+k})'| > K^k |z_{n+k}|/|z_n|$ . We apply this with  $n = 0$  and  $k = n_j$  and we may certainly assume  $|z_{n_j}|/|z_0| > 1$ . In the annular domains we also have  $|f^k(z_{n_j})'| < K'^k$ .

We may assume  $|z_{n_j}| > |z_{n_j-1}|$  so that the contraction over this part of the orbit is at least  $1/K^m$ . Then by the Koebe theorems  $B_m \subset D(z_0, M_{p_m}/K^m)$ , its distortion is bounded independent of  $m$  and its diameter goes to zero as  $m$  goes to infinity. Thus applying (5) to  $D_m$  and pulling back we obtain (2).

Thus we have shown that (2) holds at any point in  $m(I_\infty)$  and hence that its Lebesgue measure is zero.  $\square$

### 3. The results

**3.1. Two lemmas.** For a given  $f$  define the sets

$$\begin{aligned}
L &= L_f = \{z : f^n(z) \rightarrow \omega_f\}, \\
\mathcal{H}_\varepsilon &= \mathcal{H}_\varepsilon(f) = \{z : \text{dist}(z, \omega_f)\} < \varepsilon
\end{aligned}$$

where we omit the  $f$  unless confusion results.

If  $\omega_f$  is a compact repeller it is obvious that it contains no critical points. In fact, the post-singular set does not accumulate on  $\omega_f$  but actually lands on it. Precisely,

**Lemma 3.1.** *If  $\omega_f$  is a compact repeller then there is an  $\varepsilon > 0$  and an integer  $N > 0$  such that if  $c \in S$ ,  $n > N$  and  $f^n(c) \in \mathcal{K}_{\varepsilon/2}$  then  $f^n(c) \in \omega_f$ .*

*Proof.* If  $PS$  is finite, the lemma is obviously true so assume  $PS$  is not finite. Since  $\omega_f$  is compact and  $PS$  is infinite,  $\omega_f$  contains no prepoles. Moreover, since it is a repeller, there exist constants  $k > 1$  and  $\varepsilon > 0$  such that  $|f'(z)| \geq k > 1$  on the sets  $\mathcal{K}_\varepsilon$  and  $\mathcal{K}_{\varepsilon/2}$ .

Set  $V = f(\overline{\mathcal{K}_{\varepsilon/2}})$ . Since  $\overline{\mathcal{K}_{\varepsilon/2}}$  contains no poles  $V$  is compact. Moreover, since  $f|_{\mathcal{K}_{\varepsilon/2}}$  is expanding,  $\overline{\mathcal{K}_{\varepsilon/2}} \subset \text{int}(V)$  and there is a regular branch  $g$  of  $f^{-1}$  that maps  $V$  to  $\overline{\mathcal{K}_{\varepsilon/2}}$ . Thus, the annuli  $A_0 = V - g(V)$ ,  $A_{n+1} = g^n(A_0)$ ,  $n \in \mathbf{N}$  are nested and  $\text{mod}(A_n) \neq 0$  for all  $n \in \mathbf{N}$ .

We claim that for each  $n \in \mathbf{N}$

$$f^n(S) \cap (\overline{\mathcal{K}_\varepsilon} - \omega_f) = \emptyset.$$

If not, there is a sequence  $n_k \in \mathbf{N}$ ,  $n_k \rightarrow \infty$ , such that for each  $n_k$ , there is some  $c_k \in S$  (not necessarily distinct) satisfying  $v_k = f^{n_k}(c_k) \in \overline{\mathcal{K}_\varepsilon} - \omega_f$ . By the compactness of  $\overline{\mathcal{K}_\varepsilon}$ , as  $k \rightarrow \infty$ , the sequence  $v_k$  accumulates and by definition, any accumulation point belongs to  $\omega_f$ . For each  $k$ , there is an  $i(k)$  such that  $v_k \in A_{i(k)}$  and  $w_k = f^{i(k)}(v_k) \in A_0$ . Since  $\overline{A_0}$  is compact, the sequence  $w_k$  has an accumulation point in  $\overline{A_0}$  which by definition, also belongs to  $\omega_f$ . This is a contradiction because  $A_0$  is separated from  $\omega_f$  by the annuli  $A_n$ ,  $n > 0$  and  $PS \cap (\overline{\mathcal{K}} - \omega_f) = \emptyset$  as required.  $\square$

Next, for the set  $L_f$  we have

**Lemma 3.2.** *If  $f \in \mathcal{B}$  and  $\omega_f$  is a compact repeller then the Lebesgue measure of  $L_f$  is 0.*

*Proof.* Since  $\omega_f$  is a compact repeller, it is clear by the classification of stable domains that  $L_f \subset J_f$ .

Let  $\varepsilon$  be chosen as in Lemma 3.1 and define

$$\mathcal{L} = \bigcap_{n \geq 0} f^{-n}(\mathcal{K}_{\varepsilon/2}).$$

Then  $z \in \mathcal{L} - \mathcal{P}_f$  if its full forward trajectory belongs to  $\mathcal{K}_{\varepsilon/2}$ . We will prove that  $m(J_f \cap (\mathcal{L} - \mathcal{P}_f)) = 0$ . Since  $L \subset \bigcup_{n=0}^{\infty} f^{-n}(\mathcal{L})$  and  $\mathcal{P}_f$  is countable, this will imply  $m(L) = 0$ .

Suppose that  $m(\mathcal{L} - \mathcal{P}_f) > 0$  and let  $z_0$  be a density point of  $\mathcal{L} - \mathcal{P}_f$ . Since  $\omega_f$  is compact, the orbit  $\{f^n(z_0)\}$  has a finite accumulation point  $y_0 \in \overline{\mathcal{K}_{\varepsilon/2}} \subset$

$\mathcal{H}_\varepsilon$ . It follows that there exists a sequence  $n_k \rightarrow \infty$  such that  $z_k = f^{n_k}(z_0) \rightarrow y_0$ . For  $k \in \mathbf{N}$ , let  $D_k = D(z_k, \varepsilon/4)$  and let  $g_k$  be the branch of  $f^{-n_k}$  that maps  $z_k$  to  $z_0$ . Since  $z_k \in \mathcal{H}_{\varepsilon/2}$ ,  $D_k \subset \mathcal{H}_\varepsilon$  and by Lemma 3.1  $g_k$  is univalent on  $D_k$ .

Now by the definition of a density point and by Propositions 2.1 and 2.2

$$\frac{m(g_k(D_k) \cap \mathcal{L})}{m(g_k(D_k))} \rightarrow 1.$$

Again by Proposition 2.2

$$\frac{m(D_k \cap f^{n_k}(\mathcal{L}))}{m(D_k)} \rightarrow 1$$

and  $m(D_k \cap f^{n_k}(\mathcal{L})) = m(D_k)$ .

Let  $U$  be an open set with compact closure contained in  $\mathbf{C} - \overline{\mathcal{H}}$ . Since  $z_k \in J_f$  there exists an integer  $N$  such that  $f^N(D_k) \supset \bar{U}$  so that  $m(f^{N+n_k}(\mathcal{L}) \cap U) > 0$ . By definition however, for all  $k \in \mathbf{N}$ ,  $f^k(\mathcal{L}) \subset \mathcal{H}_{\varepsilon/2}$  so  $f^{N+n_k}(\mathcal{L} \cap U) = \emptyset$ . This contradiction finishes the proof.  $\square$

**3.2. The main theorem.** We are now ready to prove the main theorem.

**Theorem 3.3.** *Suppose  $f \in \mathcal{B}$ . If  $\omega_f$  is a compact repeller and if conditions (\*) and (\*\*) hold then  $f$  is ergodic with respect to Lebesgue measure on its Julia set.*

*Proof.* Suppose that  $E \subset J_f$  is an  $f$ -invariant measurable set of positive measure and let  $z$  be a density point of  $E$ . Since  $\mathcal{P}_f$  is countable we may assume  $z \notin \mathcal{P}_f$ . Under the hypotheses that  $f \in \mathcal{B}$  and that both (\*) and (\*\*) hold, by Proposition 2.3 we may assume  $z \notin I_\infty$  and thus that its orbit has a finite accumulation point  $y$ ; let  $z_k = f^{m_k}(z) \rightarrow y$ ,  $k \rightarrow \infty$ .

Under the hypothesis that  $\omega_f$  is a compact repeller, by Lemma 3.2 we may assume  $z \notin L$  and hence  $y \notin \omega_f$ . Now suppose that either  $y = f^n(v)$  for some  $v \in S$  or that for some sequence  $c_k \in S$  and iterates  $n_k$ ,  $f^{n_k}(c_k) \rightarrow y$ . By Lemma 3.1, in the first case  $f^N(y) \in \omega_f$  and in the second, the  $n_k$  are bounded and  $f^{n_k+N}(c_k) \in \omega_f$ . In either case there is an  $N$  such that  $f^N(z_k) \rightarrow \omega_f$ . Arguing as in the proof of Lemma 3.1, we can find a sequence  $k_j \rightarrow \infty$  such that  $w_j = f^{N+k_j}(z_k)$  lie in a compact annulus  $A_0$  separated from  $\omega_f$ ; since  $f^{N+k_j}(y) \in \omega_f$ , we have a contradiction. Thus  $\eta = \text{dist}(y, PS) > 0$  and we can define a univalent branch  $g_k$  of  $f^{-m_k}$  on  $D_k = D(z_k, \frac{1}{4}\eta)$  such that  $g_k(z_k) = z$ .

As above, by the definition of a density point and by Propositions 2.1 and 2.2 we obtain

$$\frac{m(g_k(D_k) \cap E)}{m(g_k(D_k))} \rightarrow 1 \quad \text{and} \quad \frac{m(D_k \cap f^{m_k}(E))}{m(D_k)} \rightarrow 1.$$

Since  $E$  is forward invariant we have  $m(D_k \cap E) = m(D_k)$ .

If  $f$  is not ergodic, we can find another forward invariant positive measure set  $F \subset J_f$ ,  $m(E \cap F) = 0$ , and a density point  $z'$  for  $F$  for which we can construct disks  $D'_l$  of fixed radius such that  $m(F \cap D'_{k'}) = m(D'_{k'})$ . Because  $z_k \in D_k$ ,  $z_{k'} \in D'_{k'}$  defined as above belong to  $J_f$  there is some  $N \in \mathbf{N}$  such that  $f^N(D'_{k'}) \supset D_k$ . It follows that  $m(E \cap F) > m(f^N(D'_{k'} \cap F) \cap (D_k \cap E)) > 0$  and we obtain a contradiction.  $\square$

#### 4. Examples

**1.** Examples of functions that the main theorem applies to are found in the class of meromorphic functions with polynomial Schwarzian derivative. By Nevanlinna's theorem, if the Schwarzian of  $f$  is a polynomial of degree  $p \geq 0$  then  $f \in \mathcal{M}_{p+2}$  and all singularities of  $f^{-1}$  are logarithmic. Moreover, at least half of the singularities are finite so it is not hard to show directly that condition  $(**)$  holds.

Now note from [19] that if the Schwarzian derivative is polynomial of degree  $p - 2$ , then in a sector of width  $2\pi/p - 2\varepsilon$  about each of the  $p$  Julia directions, the function has the asymptotic form

$$\frac{AG_\nu + BG_{\nu+1}}{CG_\nu + DG_{\nu+1}}$$

where

$$G_\nu \approx \exp(-1)^{\nu+1} z^{p/2}.$$

Therefore in any given Julia direction the coefficients  $c_p$  and functions  $\phi_p$  are all approximately equal and condition  $(*)$  holds.

The dynamical properties of these functions were described in [9]. In particular, it is shown that they have no wandering domains and no Baker domains. Thus, if each singular value lands on a repelling cycle,  $J = \widehat{\mathbf{C}}$  and the function is ergodic.

A special subclass contains the functions with constant Schwarzian derivative. To this class belong the families  $\lambda e^z$ ,  $\lambda \tan(z)$ ,  $\lambda \in \mathbf{C}$  where it is easy to construct examples for which the singular values land on repelling cycles. For instance: if  $f_1(z) = \pi i \tan(z)$ , then  $S(f_1) = \{\pm\pi\}$ ,  $PS = \{\pm\pi, 0\}$ ,  $\omega_f = \{0\}$  and  $|f'_1(0)| = \pi$ ; and if  $f_2(z) = \pi i \exp(z)$  then  $S = \{0\}$ ,  $PS = \{0, \pi i, -\pi i\}$ ,  $\omega_f = \{-\pi i\}$  and  $|f'_2(-\pi i)| = \pi$ .

**2.** Functions in the above class where Theorem 3.3 does not apply are  $e^z$  and  $\frac{1}{2}\pi i \tan(z)$ . For  $e^z$  the orbit of the singular value 0 tends to  $\infty$  and for  $\frac{1}{2}\pi i \tan(z)$  the singular values are the poles  $\pm\frac{1}{2}\pi$ ; thus, although  $\omega_f$  is a repeller, it is not compact. In fact, it is proved in [15] that  $e^z$  is not ergodic. The same is probably true for  $\pi i/2 \tan(z)$ .

Another simple example where Theorem 3.3 does not apply is the family  $a \sin(z) + b$  studied in [17]. Although the singular set consists of two points, and  $a, b$  can be chosen so that the critical values are mapped to repelling periodic orbits, condition  $(**)$  fails and  $J_f = \overline{I_\infty(f)} = \widehat{C}$  has many ergodic components.

**3.** It is hard to find examples of maps with infinitely many singular values satisfying the hypotheses of our theorem because it is hard to tell whether  $\omega_f$  is a compact repeller and to check the conditions on the Laurent expansions. Here is one such example.

Consider the entire function in  $f_\lambda = \lambda e^{-z^2} \sin(z)$ . Since it is entire condition  $(*)$  is vacuous. To see that it satisfies condition  $(**)$  note that for  $\lambda$  real, if  $|\arctan z| < \frac{1}{2}$  then  $|f'_\lambda(z)|$  is bounded. Its infinitely many critical values are bounded and accumulate on the asymptotic value 0 hence it belongs to  $\mathcal{B}$ . Moreover,  $\lambda$  may be chosen so that  $\omega_{f_\lambda}$  is a compact repeller. Thus Theorem 3.3 applies so that  $f_\lambda$  is ergodic on its Julia set.

In this case we also know there are no eventually periodic domains and, since  $m(I_\infty) = 0$ , no Baker domains. There are also no wandering domains. If there were a multiply connected wandering domain  $D$ , it would contain a homotopically non-trivial loop  $\gamma$  and by [2] the winding number of (a subsequence of)  $f^n(\gamma)$  would be non-zero for  $n$  sufficiently large and the diameter of  $f^n(\gamma)$  would tend to infinity. Thus  $\gamma$  would have to intersect the real axis; since the positive and negative real axes are asymptotic curves such domains cannot exist. Since  $m(I_\infty) = 0$ , no wandering domain can escape to infinity. Moreover, since by [7] all limit functions of  $f^n|_D$  would have to be constants in the compact repeller  $\omega_{f_\lambda}$ , we conclude  $J_{f_\lambda} = \widehat{C}$ .

### References

- [1] BAKER, I.N.: Repulsive fix points of entire functions. - Math. Z. 104, 252–256, 1968.
- [2] BAKER, I.N.: Wandering domains in the iterates of entire functions. - Proc. London Math. Soc. 49, 1984, 563–576.
- [3] BAKER, I.N., J. KOTUS, and Y. LÜ: Iterates of meromorphic functions I. - Ergodic Theory Dynam. Systems, 11, 1991, 241–248.
- [4] BAKER, I.N., J. KOTUS, and Y. LÜ: Iterates of meromorphic functions III: Preperiodic domains. - Ergodic Theory Dynam. Systems, 11, 1991, 603–618.
- [5] BAKER, I.N., J. KOTUS, and Y. LÜ: Iterates of meromorphic functions IV: Critically finite functions. - Res. Math. 22, 1992, 651–656.
- [6] BERGWEILER, W.: Iteration of meromorphic functions. - Bull. Amer. Math. Soc. 29:2, 1993, 151–188.
- [7] BERGWEILER, W., M. HARUTA, H. KRIETE, H.-G. MEIER, and N. TERGLANE: On the limit functions of iterates in wandering domains. - Ann. Acad. Fenn. Ser. A I Math. 18, 1993, 369–375.
- [8] BOCK, H.: On the dynamics of entire functions on the Julia set. - Res. Math. 30, 1996, 16–20.

- [9] DEVANEY, R.L., and L. KEEN: Dynamics of meromorphic maps with polynomial Schwarzian derivative. - *Ann. Sci. École Norm. Sup.* 22, 1989, 55–81.
- [10] EREMENKO, A., and M. LYUBICH: Iterates of entire functions. - *Soviet Math. Dokl.* 279, 1984, 25–37.
- [11] EREMENKO, A., and M. LYUBICH: Dynamical properties of some classes of entire functions. - *Ann. Inst. H. Poincaré* 42, 1992, 989–1020.
- [12] GOLDBERG, L., and L. KEEN: A finiteness theorem for a dynamical class of entire functions. - *Ergodic Theory Dynam. Systems*, 6, 1986, 183–192.
- [13] GRZEGORCZYK, P., F. PRZYTYCKI, and W. SZLENK: On the iteration of Misiurewicz's rational maps on the Riemann sphere. - *Ann. Inst. H. Poincaré* 53, 1990, 31–44.
- [14] KEEN, L., and J. KOTUS: The dynamics of the family  $\lambda \tan z$ . - *Conf. Geom. and Dyn.* 1, 1997.
- [15] LYUBICH, M.: The measurable dynamics of the exponential map. - *Siberian J. Math.* 28, 1987, 111–127.
- [16] LYUBICH, M.: On typical behavior of the trajectories of a rational mapping of the sphere. - *Soviet. Math. Dokl.* 27, 1983, 22–25.
- [17] MCMULLEN, C.: Area and Hausdorff dimension of Julia sets of entire functions. - *Trans. Amer. Math. Soc.* 300, 1987, 329–334.
- [18] MCMULLEN, C.: *Complex Dynamics and Renormalization*. - Princeton University Press, 1994.
- [19] NEVANLINNA, R.: Über Riemannsche Flächen mit endlich vielen Windungspunkten. - *Acta Math.* 58, 1932.
- [20] NEVANLINNA, R.: *Analytic Functions*. - Springer-Verlag, 1970.
- [21] RIPON, P.J., and G.M. STALLARD: Iteration of a class of hyperbolic meromorphic functions. - *Proc. Amer. Math. Soc.* (to appear).

Received 4 July 1997

Received in revised form 30 November 1998