

## ON FACTORIZATIONS OF ENTIRE FUNCTIONS OF BOUNDED TYPE

Liang-Wen Liao and Chung-Chun Yang

Nanjing University, Department of Mathematics  
Nanjing, China; maliao@nju.edu.cn

The Hong Kong University of Science & Technology  
Department of Mathematics Kowloon, Hong Kong; mayang@ust.hk

**Abstract.** We prove that if  $f$  is a transcendental entire function and the set of all finite singularities of its inverse function  $f^{-1}$  is bounded, then  $f(z) + P(z)$  is prime for any nonconstant polynomial  $P(z)$ , unless  $f(z)$  and  $P(z)$  has a nonlinear common right factor. Particularly, it is shown that  $f(z) + az$  is prime for any constant  $a \neq 0$ .

### 1. Introduction

A transcendental meromorphic function  $F$  is said to be prime (pseudo-prime) if, and only if, whenever  $F = f(g)$  for some meromorphic functions  $f$  and  $g$ , either  $f$  or  $g$  must be bilinear (rational);  $F$  is called left-prime (right-prime) if every factorization of  $F$  implies that  $f$  is bilinear whenever  $g$  is transcendental ( $g$  is linear if  $f$  is transcendental). It is easily seen  $F$  is prime if and only if  $F$  is left-prime as well as right-prime. We refer the readers to [3] or [4] for an introduction to the factorization theory of entire and meromorphic functions.

A point  $a$  is called a singularity of  $f^{-1}$  (the inverse function of  $f$ ), if  $a$  is either a critical value or asymptotic value of  $f$ . We denote by  $\text{sing}(f^{-1})$  the set of all finite singularities of  $f^{-1}$ , i.e.

$$\text{sing}(f^{-1}) = \{z \in \mathbf{C} : z \text{ is a singularity of } f^{-1}\}.$$

We denote by  $B$  the class of all entire functions  $f$  such that  $\text{sing}(f^{-1})$  is bounded and by  $S$  the class of all entire functions  $f$  such that  $\text{sing}(f^{-1})$  is finite. If  $f \in B$  ( $f \in S$ ), we say  $f$  is of bounded (finite) type.

In 1981, Noda [8] proved the following result.

**Theorem A.** *Let  $f(z)$  be a transcendental entire function. Then the set*

$$NP(f) = \{a \mid a \in \mathbf{C}, f(z) + az \text{ is not prime}\}$$

*is at most countable*

As a further study on the cardinality of  $NP(f)$ , which is denoted by  $|NP(f)|$ , Ozawa and Sawada [9] posed the following interesting question:

**Question.** *Is there any  $f$  for which the exceptional set  $NP(f)$  in Theorem A is really infinitely countable? Or what is the maximal cardinal number of the exceptional set  $NP(f)$ ?*

**Theorem B** (Ozawa and Sawada [9]). *Let  $G(w)$  be an entire function satisfying*

$$M(R, G(w)) \leq \exp(KR)$$

*for  $R \geq R_0 > 0$  and for some constant  $K > 0$ . Then either  $G(e^z) + az$  or  $G(e^z) + bz$  is prime if  $ab(a - b) \neq 0$ .*

This shows that the cardinality of  $NP(G(e^z))$  is at most 2 if  $M(R, G(w)) \leq \exp(KR)$  for  $R \geq R_0 > 0$  and for some constant  $K > 0$ . As a study of the above question, Liao–Yang [6] proved the following result.

**Theorem C.** *Let  $f$  be a transcendental entire function of finite order in  $S$ . Then for any constant  $a \neq 0$ ,  $f(z) + az$  is prime, i.e.  $|NP(f)| \leq 1$ .*

Recently, Wang–Yang [13] proved the following theorem.

**Theorem D.** *Let  $P, Q$  be nonconstant polynomials,  $\alpha \in B$ ,  $h$  a periodic entire function of order one and mean type,  $G(z) = P \circ h \circ \alpha(z)$ . If  $F(z) = G^n(z) + Q(z)$  has a factorization  $F(z) = f(g(z))$ , then  $g(z)$  must be a common right factor of  $\alpha(z)$  and  $Q(z)$ .*

**Remark 1.** The original statement of Theorem D only requires that  $h$  is of order one. Here we would like to point out that  $h$  should be at most order one of mean type, as it is needed in the proof of Theorem D, Lemma 5 in [13]. However,  $f$  in Lemma 5 should be an entire function of exponential type, i.e.  $f$  has order less than one or order one and mean type; see p. 27 in [4].

**Remark 2.** Let  $G$  be defined in Theorem D. Then  $G^n(z) + az$  is prime for any constant  $a \neq 0$ .

As a continuation of the study of our previous work [6], we are able to extend Theorem C to a large class of functions, namely, functions of bounded type. The following is our main result.

**Theorem.** *Let  $f$  be a transcendental entire function in  $B$ , then for any nonconstant polynomial  $P(z)$ ,  $f(z) + P(z)$  is prime unless  $f(z)$  and  $P(z)$  has a nonlinear common right factor.*

**2. Some lemmas**

**Lemma 1** (Rippon and Stallard [11]). *Let  $f$  be a meromorphic function with a bounded set of all finite critical and asymptotic values. Then there exists  $K > 0$  such that if  $|z| > K$  and  $|f(z)| > K$ , then*

$$|f'(z)| \geq \frac{|f(z)| \log |f(z)|}{16\pi|z|}.$$

**Lemma 2** ([5]). *Let  $f$  be a transcendental entire function, and  $0 < \delta < \frac{1}{4}$ . Suppose that at the point  $z$  with  $|z| = r$  the inequality*

$$(1) \quad |f(z)| > M(r, f)\nu(r, f)^{-(1/4)+\delta}$$

*holds. Then there exists a set  $F$  in  $R^+$  and of finite logarithmic measure, i.e.,*

$$\int_F \frac{dt}{t} < +\infty$$

*such that*

$$(2) \quad f^{(m)}(z) = \left(\frac{\nu(r, f)}{z}\right)^m (1 + o(1))f(z)$$

*holds whenever  $m$  is a fixed nonnegative integer and  $r \notin F$ .*

**Lemma 3** (Baker and Singh [1], also see [2]). *Let  $f$  and  $g$  be two entire functions. Then*

$$\text{sing}((f \circ g)^{-1}) \subset \text{sing}(f^{-1}) \cup f(\text{sing}(g^{-1})).$$

**Lemma 4** (Polya [10]). *Let  $f$  and  $g$  be two transcendental entire functions. Then*

$$\lim_{r \rightarrow \infty} \frac{M(r, f \circ g)}{M(r, g)} = \infty.$$

**Lemma 5.** *Let  $f$  be a transcendental entire function. Then*

$$M(r, f') \leq M(r, f)^2$$

*for a sufficiently large  $r$ .*

**Remark 3.** This follows easily from a result of Valiron ([12]):

$$\lim_{r \rightarrow \infty} \frac{\log M(r, f')}{\log M(r, f)} = 1.$$

### 3. Proof of the theorem

Let  $F(z) = f(z) + P(z)$ ,  $P(z)$  is a nonconstant polynomial. We first prove that  $F$  is pseudo-prime. Assume that

$$F(z) = g(h(z)),$$

where  $g$  is a transcendental meromorphic function with at most one pole and  $h$  is a transcendental entire function. Thus

$$(3) \quad f(z) = g(h(z)) - P(z), \quad f'(z) = g'(h(z))h'(z) - P'(z).$$

First we consider the case that  $g$  is a transcendental entire function, and then we discuss two situations.

Case 1:  $g'$  has at least two zeros. Then there exists a zero  $c$  of  $g'$  such that  $h(z) = c$  has infinitely many roots  $\{z_k\}_{k=1}^{\infty}$ . Thus we have

$$f(z_k) = -P(z_k) + g(c), \quad f'(z_k) = -P'(z_k).$$

By Lemma 1, we would have

$$|P'(z_k)| \geq \frac{|P(z_k) - g(c)| \log |P(z_k) - g(c)|}{16\pi |z_k|},$$

which leads to a contradiction.

Case 2:  $g'$  has at most one zero. Thus

$$g'(w) = (w - w_0)^n e^{\alpha(w)}, \quad f'(z) = (h(z) - w_0)^n e^{\alpha(h(z))} h'(z) - P'(z),$$

where  $n$  is a non-negative integer. Let  $K(z) = e^{-\alpha(h(z))/(n+3)}$ , and assume that  $\Gamma$  is a simple curve tending to infinity such that if  $z \in \Gamma$  and  $|z| = r$ , then  $|K(z)| = M(r, K)$ . By Lemmas 4 and 5, we have, if  $z \in \Gamma$  and  $|z| = r$  is sufficiently large,

$$(4) \quad \begin{aligned} |g'(h(z))h'(z)| &= |(h(z) - w_0)^n e^{\alpha(h(z))} h'(z)| \\ &= \frac{|(h(z) - w_0)^n h'(z)|}{M(r, K)^{n+3}} \leq \frac{1}{M(r, K)} \rightarrow 0. \end{aligned}$$

Let  $L(z) = -\alpha(h(z))/(n+3)$  and  $A(r, L) = \max_{|z|=r} \operatorname{Re} L(z)$ . Thus if  $z \in \Gamma$ ,  $|K(z)| = M(r, K) = e^{A(r, L)}$ ,  $\operatorname{Re} L(z) = A(r, L)$ . By Hadamard's three-circle theorem, we have, for  $r_1 < r_2 < r_3$ ,

$$(5) \quad A(r_2, L) \leq \frac{\log r_2 - \log r_1}{\log r_3 - \log r_1} A(r_3, L) + \frac{\log r_3 - \log r_2}{\log r_3 - \log r_1} A(r_1, L).$$

For  $z_0 \in \Gamma$ , we have

$$(6) \quad |L'(z_0)| = \lim_{z \rightarrow z_0, z \in \Gamma} \frac{|L(z) - L(z_0)|}{|z - z_0|} \geq \lim_{z \rightarrow z_0, z \in \Gamma} \frac{|\operatorname{Re} L(z) - \operatorname{Re} L(z_0)|}{|z - z_0|}.$$

Let  $|z_0| = r_0$  and  $|z| = r_0 + h$ ,  $h > 0$ , then as  $z \rightarrow z_0$ ,  $h \rightarrow 0$ . Thus, by (5) and (6), we have, for sufficiently large  $r_0$ ,

$$(7) \quad \begin{aligned} |L'(z_0)| &\geq \lim_{z \rightarrow z_0, z \in \Gamma} \frac{A(r_0 + h, L) - A(r_0, L)}{|z - z_0|} \\ &= \lim_{z \rightarrow z_0, z \in \Gamma} \frac{h}{|z - z_0|} \frac{A(r_0 + h, L) - A(r_0, L)}{h} \\ &= \lim_{h \rightarrow 0} \frac{A(r_0 + h, L) - A(r_0, L)}{h} \\ &\geq \lim_{h \rightarrow 0} \frac{\frac{\log(1 + h/r_0)}{\log r_0} (A(r_0, L) - A(1, L))}{h} \\ &= \frac{A(r_0, L) - A(1, L)}{r_0 \log r_0} > 1. \end{aligned}$$

Let  $w = G(z) = e^{\alpha(h(z))/(n+3)} = e^{-L(z)}$ . Thus 0 is an asymptotic value of  $G$  and  $\Gamma$  is the corresponding asymptotic curve,  $\gamma = G(\Gamma)$  is a simple curve connecting  $G(0)$  and 0. Let  $B$  be the length of  $\gamma$ , which is a finite number. And  $dw = e^{-L(z)}L'(z)dz$ . By this, (4) and (7), if  $z \in \Gamma$ , we have

$$\begin{aligned} |g(h(z))| &= \left| \int_{z_0 \text{ along } \Gamma}^z g'(h(z))h'(z) dz + g(h(z_0)) \right| \\ &\leq \int_{z_0 \text{ along } \Gamma}^z |g'(h(z))h'(z)| |dz| + |g(h(z_0))| \\ &\leq \int_{w_0 \text{ along } \gamma}^w \frac{1}{|L'(z)|} |dw| + |g(h(z_0))| \\ &\leq \int_{w_0 \text{ along } \gamma}^w |dw| + |g(h(z_0))| \\ &\leq B + |g(h(z_0))|. \end{aligned}$$

Thus we can find a sequence of  $\{z_k\}_{k=1}^\infty$  such that  $z_k \rightarrow \infty$  as  $k \rightarrow \infty$ , and

$$f(z_k) \sim -P(z_k), \quad f'(z_k) \sim -P'(z_k).$$

A contradiction follows from this and Lemma 1.

If  $g'$  has just one pole  $w_1$ , so does  $g$ , then  $h(z)$  does not assume  $w_1$ , i.e.,  $h(z) = e^{\beta(z)} + w_1$ . Moreover, if  $g'$  has a zero  $c$ , then  $h(z) = c$  has infinitely many roots. One can derive a contradiction by arguing similarly as in Case 1. Hence  $g'$  has no zeros, i.e.,

$$g'(w) = \frac{1}{(w - w_1)^n} e^{\alpha(w)},$$

and

$$g'(h(z))h'(z) = \beta'(z) \exp(\alpha(e^{\beta(z)} + w_1)) + (1 - n)\beta(z).$$

By the same argument as that in Case 2 above, we can get a contradiction. Thus  $F(z) = f(z) + P(z)$  is pseudo-prime. Now we assume that  $F(z)$  has the following factorization:

$$F(z) = f(z) + P(z) = Q(g(z)),$$

where  $Q$  is rational,  $g$  is a transcendental meromorphic function. If  $Q$  is a polynomial, then  $g$  is entire. If  $Q$  has a pole  $w_1$ , then  $g(z)$  does not assume  $w_1$ . Thus  $h(z) = 1/(g(z) - w_1)$  is an entire function and  $F(z) = Q_1(h(z))$ , where  $Q_1$  is a rational function. Without loss of generality, we may assume that  $g(z)$  is entire, and  $Q(w)$  has at most one pole. Now we discuss the following two sub-cases.

*Subcase 1:*  $Q$  has one pole, say  $w_0$ , i.e.,  $Q(w) = Q_1(w)/(w - w_0)^n$ , where  $Q_1(w)$  is a polynomial with degree  $m$  and  $Q_1(w_0) \neq 0$ . Then  $g(z) = w_0 + e^{h(z)}$ , where  $h(z)$  is a nonconstant entire function. Thus we have

$$\begin{aligned} f(z) &= Q_1(w_0 + e^{h(z)})e^{-nh(z)} - P(z) \\ &= a_0e^{-nh(z)} + a_1e^{-(n-1)h(z)} + \dots + a_me^{(m-n)h(z)} - P(z), \end{aligned}$$

where  $a_0, a_1, \dots, a_m$  are constants and  $a_m \neq 0$ ,  $a_0 = Q_1(w_0) \neq 0$ . Thus

$$\begin{aligned} f'(z) &= (-na_0e^{-nh(z)} - (n-1)a_1e^{-(n-1)h(z)} + \dots \\ &\quad + (m-n)a_me^{(m-n)h(z)})h'(z) - P'(z) \\ &= [-na_0 - (n-1)a_1e^{h(z)} + \dots \\ &\quad + (m-n)a_me^{mh(z)}]e^{-nh(z)}h'(z) - P'(z) \\ &= P_1(e^{h(z)})e^{-nh(z)}h'(z) - P'(z), \end{aligned}$$

where  $P_1(w)$  is a polynomial and  $P_1(0) = -na_0 \neq 0$ . If  $P_1(w)$  is a nonconstant polynomial, then  $P_1(w)$  has a zero  $c \neq 0$  and  $e^{h(z)} = c$  has infinitely many roots. Let  $\{z_k\}_{k=1}^{+\infty}$  be zeros of  $e^{h(z)} - c$ , then  $f'(z_k) = -P'(z_k)$  and

$$f(z_k) = \frac{Q_1(w_0 + c)}{c^n} - P(z_k).$$

Again, by Lemma 1, we have a contradiction. If  $P_1(w)$  is a constant polynomial, then

$$f(z) = a_0 e^{-nh(z)} + a_m - P(z), \quad f'(z) = -na_0 e^{-nh(z)} h'(z) - P'(z).$$

Let  $K(z) = e^{nh(z)}$  and  $|z'| = r$ ,  $|K(z')| = M(r, K)$ . Then by Lemma 2, we have, for  $r \notin F$ ,

$$\begin{aligned} |-na_0 e^{-nh(z')} h'(z')| &= \left| a_0 \frac{1}{K(z')} \frac{K'(z')}{K(z')} \right| = |a_0| \frac{1}{M(r, K)} \frac{\nu(r, K)}{r} (1 + o(1)), \\ |a_0 e^{-nh(z')}| &= \frac{|a_0|}{M(r, K)}. \end{aligned}$$

Noting  $\lim_{r \rightarrow \infty} (\nu(r, K)/M(r, K)) = 0$  for a transcendental entire function  $K$ , we can find a sequence of  $\{z_k\}_{k=1}^{+\infty}$  such that  $|f(z_k)| \sim |P(z_k)|$ ,  $|f'(z_k)| \sim |P'(z_k)|$ . A contradiction follows from this and Lemma 1.

*Subcase 2:*  $Q(w)$  has no pole, i.e.,  $Q(w)$  is a polynomial with degree  $\geq 2$ . If  $Q'(w)$  has at least two distinct zeros, then there exists a zero  $w_1$  of  $Q'(w)$  such that  $g(z) = w_1$  has infinitely many zeros  $\{z_n\}_{n=1}^{+\infty}$ . Then

$$f'(z_n) = Q'(g(z_n)) - P'(z_n) = -P'(z_n), \quad f(z_n) = Q(w_1) + P(z_n).$$

However, by Lemma 1,

$$|f'(z_n)| \geq \frac{|f(z_n)| \log |f(z_n)|}{16\pi |z_n|},$$

which will lead to a contradiction. Therefore, we only need to treat the case that  $Q'(w)$  has only one zero  $w_0$ . If  $g(z) - w_0$  has infinitely many zeros, again a contradiction follows from Lemma 1. Hence, we have

$$g(z) = w_0 + p_1(z)e^{h(z)} \quad \text{and} \quad Q'(z) = A(w - w_0)^{n-1},$$

where  $p_1(z)$  is a polynomial,  $h(z)$  a nonconstant entire function. Thus

$$\begin{aligned} Q(w) &= \frac{A}{n}(w - w_0)^n + B, \\ f(z) &= \frac{A}{n} p_1(z)^n e^{nh(z)} + B - P(z), \\ f'(z) &= \frac{A}{n} (p_1'(z) + p_1(z)nh'(z)) e^{nh(z)} - P'(z). \end{aligned}$$

Set  $K(z) = e^{-nh(z)}$  and let  $|z'| = r$ ,  $K(z') = M(r, K)$ . Then it follows from Lemma 2, for  $r \notin F$ , that

$$\begin{aligned} \left| \frac{A}{n} (p_1'(z') + p_1(z')nh'(z'))e^{nh(z')} \right| &= \left| \frac{A}{n} \left( \frac{p_1'(z')}{K(z')} - \frac{p_1(z')}{K(z')} \frac{K'(z')}{K(z')} \right) \right| \\ &\leq \frac{cr^t}{M(r, K)} + \frac{dr^t\nu(r, K)}{M(r, K)}, \end{aligned}$$

where  $c, d$  are positive constants,  $t = \deg p_1 - 1$ . Noting

$$\lim_{r \rightarrow \infty} \frac{r^t\nu(r, K)}{M(r, K)} = 0$$

for a transcendental entire function  $K$ , there exists a sequence of  $\{z_n\}_{n=1}^{+\infty}$  such that

$$f(z_n) \sim -P(z_n), \quad f'(z_n) \sim -P'(z_n).$$

Again by Lemma 1, we get a contradiction. Thus we have proved that  $F(z) = f(z) + P(z)$  is left-prime. Next we show that  $F$  is right-prime. Let

$$F(z) = g(q(z)),$$

where  $g$  is a transcendental entire function and  $q(z)$  a polynomial with degree  $\geq 2$ . Thus

$$f(z) = g(q(z)) - P(z)$$

and hence

$$f'(z) = g'(q(z))q'(z) - P'(z).$$

First, we prove that  $g'(w)$  has infinitely many zeros. In fact, if  $g'(w)$  has only finitely many zeros, then  $g'(w) = s(w)e^{h(w)}$ , where  $s(w)$  is a polynomial and  $h(w)$  is a nonconstant entire function. Let  $K(z) = e^{-h(z)/3}$ . There exists a curve  $\Gamma$  tending to infinity such that if  $z \in \Gamma$ , then  $|K(z)| = M(|z|, K)$ . Noting that  $K$  is a transcendental entire function, we have that  $M(r, K) \geq r^{2m+2}$  for  $r \geq r_0$ , where  $m = \deg s$ . Let  $w = G(z) = e^{h(z)/3}$  and  $\lambda = G(\Gamma)$ . Then  $dw = \frac{1}{3}h'(z)e^{h(z)/3}$ . If  $h(z)$  is nonconstant polynomial, then there exists a positive constant  $c$  such that  $|h'(z)| \geq c$  for sufficiently large  $|z| = r$ . If  $h(z)$  is transcendental, then  $|\frac{1}{3}h'(z)| > 1$  for  $z \in \Gamma$  and sufficiently large  $|z| = r$ , by (7). Hence, we have, for  $z \in \Gamma$  and  $|z| \geq r_0$ ,

$$\begin{aligned} |g'(z)| &\leq \frac{1}{M(r, K)^2}, \\ |g(z)| &= \left| \int_{z_0 \text{ along } \Gamma}^z g'(z) dz + g(z_0) \right| \leq \left| \int_{w_0 \text{ along } \lambda}^w |dw| \right| \leq A, \end{aligned}$$

where  $w_0 = G(z_0)$ ,  $w = G(z)$  and  $A$  is a positive constant. Let  $\gamma$  be a component of  $q^{-1}(\Gamma)$ , and denote  $R = |q(z)|$  for  $z \in \gamma$ . Then for  $z \in \gamma$ , we have

$$|g(q(z))| \leq A, \quad |g'(z)q'(z)| \leq \frac{BR^{m+1}}{M(R, K)^2} \rightarrow 0, \quad \text{as } z \rightarrow \infty,$$

where  $A$  and  $B$  are constants. Hence, for  $z \in \gamma$ , we have

$$|f(z)| \sim |P(z)|, \quad |f'(z)| \sim |P'(z)|.$$

Again, by Lemma 1, the above estimates will lead to a contradiction as before. Thus  $g'$  has infinitely many zeros. Now let  $n = \deg q$  and  $m = \deg P$ . Next we will prove that  $n \mid m$ , i.e., there is a positive integer  $r$  such that  $m = nr$ . Let  $\{w_k\}_{k=1}^\infty$  denote the zeros of  $g'(w)$  and set

$$q(z) = a_n z^n + a_{n-1} z^{n-1} + \dots + a_1 z + a_0.$$

We consider the roots of the equation

$$q(z) = w_k,$$

which implies

$$(8) \quad a_n z^n (1 + o(1)) = w_k.$$

On the other hand, the roots of the above equation can be expressed as

$$z_k^{(j)} = \left| \frac{w_k}{a_n} \right|^{1/n} e^{i(2j\pi + \phi_k)/n} (1 + o(1)),$$

where

$$\phi_k = \arg \frac{w_k}{a_n}, \quad j = 0, 1, 2, \dots, n - 1.$$

Thus

$$\begin{aligned} P(z_k^{(0)}) &\sim A|w_k|^{m/n}, \\ P(z_k^{(1)}) &\sim e^{2m\pi i/n} A|w_k|^{m/n}, \\ P'(z_k^{(0)}) &\sim B|w_k|^{(m-1)/n}, \\ P'(z_k^{(1)}) &\sim e^{2(m-1)\pi i/n} B|w_k|^{(m-1)/n}, \end{aligned}$$

where  $A, B$  are constants depending on  $q(z)$  and  $P(z)$  only. Thus we have sequences  $\{w_k\}_{k=1}^\infty$ , with  $w_k \rightarrow \infty$  as  $k \rightarrow \infty$ ,  $\{z_k^{(0)}\}_{k=1}^\infty$  and  $\{z_k^{(1)}\}_{k=1}^\infty$  such that

- (9)  $q(z_k^{(0)}) = q(z_k^{(1)}) = w_k,$
- (10)  $P(z_k^{(0)}) - P(z_k^{(1)}) \sim (1 - e^{2m\pi i/n})A|w_k|^{m/n},$
- (11)  $f'(z_k^{(0)}) = -P'(z_k^{(0)}) \sim -B|w_k|^{(m-1)/n},$
- (12)  $f'(z_k^{(1)}) = -P'(z_k^{(1)}) \sim -e^{2(m-1)\pi i/n}B|w_k|^{(m-1)/n},$
- (13)  $f(z_k^{(0)}) = g(w_k) - P(z_k^{(0)}),$
- (14)  $f(z_k^{(1)}) = g(w_k) - P(z_k^{(1)}),$
- (15)  $f(z_k^{(1)}) - f(z_k^{(0)}) = P(z_k^{(0)}) - P(z_k^{(1)}).$

If  $n \nmid m$ , then  $1 - e^{2m\pi i/n} \neq 0$ . Now we discuss two subcases.

*Subcase 1:*  $\{f(z_k^{(0)})\}_{k=1}^\infty$  is bounded. We have, by (10)–(15),

$$(16) \quad |f(z_k^{(1)})| \sim |(1 - e^{2m\pi i/n})A| |w_k|^{m/n}.$$

By this and Lemma 1, we obtain that

$$\begin{aligned} |B| |w_k|^{(m-1)/n} \sim |f'(z_k^{(1)})| &\geq \frac{|f(z_k^{(1)})| \log |f(z_k^{(1)})|}{16\pi |z_k^{(1)}|} \\ &\sim C |w_k|^{(m-1)/n} \log(|(1 - e^{2m\pi i/n})A| |w_k|^{m/n}), \end{aligned}$$

where

$$C = \frac{|(1 - e^{2m\pi i/n})A| |a_n|^{1/n}}{16\pi},$$

which is a contradiction.

*Subcase 2:*  $\{f(z_k^{(0)})\}_{k=1}^\infty$  is unbounded. Then there exists a sub-sequence of  $\{f(z_k^{(0)})\}_{k=1}^\infty$  tending to infinity, which we may, without confusing, denote by the original sequence:  $\{f(z_k^{(0)})\}_{k=1}^\infty$ . Thus by Lemma 1, we have

$$\begin{aligned} |B| |w_k|^{(m-1)/n} \sim |f'(z_k^{(0)})| &\geq \frac{|f(z_k^{(0)})| \log |f(z_k^{(0)})|}{16\pi |z_k^{(0)}|} \\ &\sim \frac{|a_n|^{1/n} |f(z_k^{(0)})| \log |f(z_k^{(0)})|}{16\pi |w_k|^{1/n}}. \end{aligned}$$

Hence,

$$|f(z_k^{(0)})| = o(|w_k^{(m/n)}|).$$

Thus

$$|f(z_k^{(1)})| \sim |(1 - e^{2m\pi i/n})A| |w_k^{m/n}|.$$

By arguing similarly as in Subcase 1, we will arrive at a contradiction. Hence  $n \mid m$ . Finally, we will prove that  $q(z)$  is a common right factor of  $f(z)$  and  $P(z)$ . If  $q(z)$  is not a right factor of  $P(z)$ , then there exist polynomials  $Q$  and  $P_1$  with  $0 < \deg P_1 < n = \deg q$  such that

$$P(z) = Q(q(z)) + P_1(z).$$

Thus

$$G(z) = f(z) + P_1(z) = g(q(z)) - Q(q(z)) = g_1(q(z)),$$

where  $g_1(w) = g(w) - Q(w)$  is a transcendental entire function. By arguing similarly as in the subcase above, it follows that  $n \mid \deg P_1$ , which is a contradiction. Thus,  $P(z) = Q(q(z))$  and  $f(z) = g(q(z)) - Q(q(z))$ . The conclusion follows.

#### 4. Concluding remarks

**Corollary.** *Let  $f$  be a transcendental entire function in  $B$ , then for any constant  $a \neq 0$ ,  $f(z) + az$  is prime.*

**Remark 4.** This corollary shows that if  $f(z) - az \in B$  for some constant  $a$ , then  $|NP(f)| \leq 1$ .

**Remark 5.** If  $h$  is a periodic entire function of order one and mean type, then  $h \in B$ . Thus if  $G(z)$  is as stated in Theorem D, then  $G^n \in B$ .

**Remark 6.** The condition  $f \in B$  in the above theorem and corollary is not removable. For example,  $f(z) = e^z e^{e^z} + e^z$ , then  $f(z) = (we^w + w) \circ e^z$ , and  $f(z) + z = (e^w + w) \circ (e^z + z)$ . This example shows the cardinality of  $NP(f)$  may be greater than one if  $f \notin B$ .

**Remark 7.** If  $f$  is an entire function such that  $\text{sing}(f^{-1}) \subset \mathbf{R}$ , then, by Lemma 3,  $\sin(f(z)) \in B$  and  $\cos(f(z)) \in B$ . Thus, for any constant  $a \neq 0$ ,  $\sin(f(z)) + az$  and  $\cos(f(z)) + az$  are prime. It was mentioned in [2] that the Pólya–Laguerre class  $LP$  consists of all entire functions  $f$  which have a representation

$$f(z) = \exp(-az^2 + bz + c)z^n \prod \left(1 - \frac{z}{z_k}\right) \exp\left(\frac{z}{z_k}\right),$$

where  $a, b, c \in \mathbf{R}$ ,  $a \geq 0$ ,  $n \in \mathbf{N}_0$ ,  $z_k \in \mathbf{R} \setminus \{0\}$  for all  $k \in \mathbf{N}$ , and  $\sum_{k=1}^\infty |z_k|^{-2} < \infty$ . Furthermore, if  $f_1, f_2, \dots, f_n \in LP$ , and  $f = f_1 \circ f_2 \circ \dots \circ f_n$ , then  $\text{sing}(f^{-1}) \subset \mathbf{R}$ . Thus, for example,  $\sin(f(z)) + az$  is prime for  $a \neq 0$ , when  $f \in LP$ .

**References**

- [1] BAKER, I. N., and A. SINGH: Wandering domains in the iteration of compositions of entire functions. - *Ann. Acad. Sci. Fenn. Math.* 20, 1995, 149–153.
- [2] BERGWELER, W., and Y. F. WANG: On the dynamics of composite entire functions. - *Ark. Mat.* 36, 1998, 31–39.
- [3] CHUANG, CHI-TAI, and CHUANG-CHUN YANG: Fix-Points and Factorization of Meromorphic Functions. - World Scientific, Singapore–New Jersey–London–Hong Kong, 1990.
- [4] GROSS, F.: Factorization of Meromorphic Functions. - U. S. Government Printing Office, 1972.
- [5] LAINE, I.: Nevanlinna Theory and Complex Differential Equations. - Walter de Gruyter, Berlin–New York, 1993.
- [6] LIAO, L. W., and C. C. YANG: On factorizations of entire functions of finite type. - Preprint.
- [7] NEVANLINNA, R.: Eindeutige analytische Funktionen. - Springer-Verlag, Berlin–Göttingen–Heidelberg, 1953.
- [8] NODA, Y.: On factorization of entire functions. - *Kodai Math. J.* 4, 1981, 480–494.
- [9] OZAWA, M., and K. SAWADA: A theorem on factorization of certain entire functions. - *Complex Variables Theory Appl.* 34, 1997, 181–185.
- [10] POLYA, G.: On an integral of an integral function. - *J. London Math. Soc.* 1, 1926, 12–15.
- [11] RIPPON, P. J., and G. M. STALLARD: Iteration of a class of hyperbolic meromorphic functions. - *Proc. Amer. Math. Soc.* 127, 1999, 3251–3258.
- [12] VALIRON, G.: Lecture on the general theory of integral functions. -Toulouse, Edouard privat, 1923.
- [13] WANG, X. L., and C. C. YANG: On the factorization of a certain class of entire functions. - *Indian J. Pure Appl. Math.* 33, 2002, 1727–1741.

Received 1 October 2003